



JC11 Rec'd PCT/PTO SEP 26 2001

FORM PTO-1390  
(REV. 11-2000)

DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE

ATTORNEY'S DOCKET NUMBER

TRANSMITTAL LETTER TO THE UNITED STATES  
DESIGNATED/ELECTED OFFICE (DO/EO/US)  
CONCERNING A FILING UNDER 35 U.S.C. 371

1422-0493P

U.S. APPLICATION NO. 09/937521  
NEW

INTERNATIONAL APPLICATION NO.

INTERNATIONAL FILING DATE

PRIORITY DATE CLAIMED

PCT/JP00/01802

March 24, 2000

March 26, 1999

TITLE OF INVENTION

CERAMIDASE GENE

APPLICANT(S) FOR DO/EO/US

ITO, Makoto

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☒ This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39 (1).
4. ☐ The US has been elected by the expiration of 19 months from the priority date (Article 31).
5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2))
  - a. ☐ is transmitted herewith (required only if not transmitted by the International Bureau).
  - b. ☒ has been transmitted by the International Bureau. WO 00/58448
  - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☒ An English language translation of the International Application as filed (35 U.S.C. 371(c)(2)).
  - a. ☒ is transmitted herewith.
  - b. ☐ has been previously submitted under 35 U.S.C. 154(d)(4)
7. ☒ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3)).
  - a. ☐ are transmitted herewith (required only if not transmitted by the International Bureau).
  - b. ☐ have been transmitted by the International Bureau.
  - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
  - d. ☒ have not been made and will not be made.
8. ☐ An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
9. ☒ An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).
10. ☐ An English language translation of the annexes of the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

Items 11. to 20. below concern document(s) or information included:

11. ☒ An Information Disclosure Statement under 37 CFR 1.97 and 1.98, Form PTO-1449(s), and International Search Report (PCT/ISA/210) with 8 cited document(s).
12. ☒ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
13. ☒ A **FIRST** preliminary amendment.
14. ☐ A **SECOND** or **SUBSEQUENT** preliminary amendment.
15. ☐ A substitute specification.
16. ☐ A change of power of attorney and/or address letter.
17. ☐ A computer-readable form of the sequence listing in accordance with PCT Rule 13ter.2 and 35 U.S.C. 1.821-1.825.
18. ☐ A second copy of the published international application under 35 U.S.C. 154(d)(4).
19. ☐ A second copy of the English language translation of the international application under 35 U.S.C. 154(d)(4).
20. ☒ Other items or information:  
International Preliminary Report (PCT/IPEA/409)  
Two (2) Sheets of Formal Drawings  
Sequence Listing

Form PTO-1390 (REV 11-2000) page 2 of 2

PATENT  
1422-0493P

IN THE U.S. PATENT AND TRADEMARK OFFICE

Applicant: ITO, Makoto Conf.:  
Int'l. Appl. No.: PCT/JP00/01802  
Appl. No.: New Group:  
Filed: September 26, 2001 Examiner:  
For: CERAMIDASE GENE

PRELIMINARY AMENDMENT

**BOX PATENT APPLICATION**

Assistant Commissioner for Patents  
Washington, DC 20231

September 26, 2001

Sir:

The following Preliminary Amendments and Remarks are respectfully submitted in connection with the above-identified application.

AMENDMENTS

IN THE SPECIFICATION:

Please amend the specification as follows:

Before line 1, insert --This application is the national phase under 35 U.S.C. § 371 of PCT International Application No. PCT/JP00/01802 which has an International filing date of March 24, 2000, which designated the United States of America and was not published in English.--

IN THE CLAIMS:

Please amend the claims as follows:

4. (Amended) A recombinant DNA comprising the gene of claim 1.

5. (Amended) An expression vector for a microorganism, an animal cell or a plant cell, comprising the gene of claim 1 or the recombinant DNA of claim 4.

9. (Amended) A polypeptide possessing a ceramidase activity, encoded by the gene of claim 1.

11. (Amended) An antisense DNA which is complementary to the gene of claim 1 or a part thereof.

12. (Amended) An Antisense RNA which is complementary to the gene of claim 1 or a part thereof.

14. (Amended) An oligonucleotide probe or primer, capable of specifically hybridizing to the gene of claim 1, or a complementary stand thereof.

15. (Amended) An antibody or a fragment thereof, capable of specifically binding to the polypeptide of claim 8.

20. (Amended) A method of controlling an amount of ceramide in a cell and/or in a tissue, characterized by introducing the gene of claim 1 or an antisense nucleic acid thereof into the cell and/or the tissue, thereby controlling the amount of ceramide in the cell and/or in the tissue.

109260 "T 054660

REMARKS

The specification has been amended to provide a cross-reference to the previously filed International Application.

The claims have also been amended to delete improper multiple dependencies.

Entry of the above amendments is earnestly solicited. An early and favorable first action on the merits is earnestly solicited.

If necessary, the Commissioner is hereby authorized in this, concurrent, and future replies, to charge payment or credit any overpayment to Deposit Account No. 02-2448 for any additional fees required under 37 C.F.R. § 1.16 or under 37 C.F.R. § 1.17; particularly, extension of time fees.

Respectfully submitted,

BIRCH, STEWART, KOLASCH & BIRCH, LLP

By 

Marc S. Weiner, #32,181

P.O. Box 747

Falls Church, VA 22040-0747

(703) 205-8000

MSW/rem

1422-0493P

Attachment: VERSION WITH MARKINGS TO SHOW CHANGES MADE

VERSION WITH MARKINGS TO SHOW CHANGES MADE

The specification has been amended to provide a cross-reference to the previously filed International Application.

IN THE CLAIMS:

The claims have been amended as follows:

4. (Amended) A recombinant DNA comprising the gene of [any one of claims 1 to 3] claim 1.

5. (Amended) An expression vector for a microorganism, an animal cell or a plant cell, comprising the gene of [any one of claims 1 to 3] claim 1 or the recombinant DNA of claim 4.

9. (Amended) A polypeptide possessing a ceramidase activity, encoded by the gene of [any one of claims 1 to 3] claim 1.

11. (Amended) An antisense DNA which is complementary to the gene of [any one of claims 1 to 3] claim 1 or a part thereof.

12. (Amended) An Antisense RNA which is complementary to the gene of [any one of claims 1 to 3] claim 1 or a part thereof.

14. (Amended) An oligonucleotide probe or primer, capable of specifically hybridizing to the gene of [any one of claims 1 to 3] claim 1, or a complementary stand thereof.

15. (Amended) An antibody or a fragment thereof, capable of specifically binding to the polypeptide of [any one of claims 8 to 10] claim 8.

20. (Amended) A method of controlling an amount of ceramide in a cell and/or in a tissue, characterized by introducing the gene of [any one of claims 1 to 3] claim 1 or an antisense nucleic acid thereof into the cell and/or the tissue, thereby controlling the amount of ceramide in the cell and/or in the tissue.

2/PATS

1

09/937521

JC12 Rec'd PCT/FTO 26 SEP 2001

## DESCRIPTION

### CERAMIDASE GENE

#### 5 TECHNICAL FIELD

10 The present invention relates to a polypeptide possessing a ceramidase activity, an antibody specifically binding thereto, a gene encoding the polypeptide, and a probe and primer which are capable of specifically hybridizing thereto, which are useful as a reagent for lipid engineering for analyzing a structure, functions, and the like of a ceramide. In addition, the present invention relates to a method for producing the above-mentioned polypeptide by a genetic engineering means, and a method for detecting the polypeptide or the gene, and a kit therefor. Further, the present invention relates to a method of controlling an amount of a ceramide in a cell and/or in a tissue, 15 which can be applied to a disease caused by abnormality in the amount of the ceramide.

#### BACKGROUND ART

20 A ceramidase is an enzyme which hydrolyzes a ceramide, a kind of a sphingolipid, into a sphingoid and a fatty acid. The sphingoid which is generated by hydrolyzing the ceramide with the ceramidase possesses various physiological activities such as inhibition of protein kinase C, activation of phospholipase D and inhibition of a calmodulin-dependent enzyme. As described above, the above-mentioned sphingoid is an important substance 25 which is thought to be acting on the regulation of the cell functions because the

sphingoid is involved in proliferation of cells and intracellular signal transduction. The ceramidase is an enzyme which plays an important role of the control of the amount of the above-mentioned sphingoid.

Ceramidases are classified into acidic ceramidases and neutral/alkaline ceramidases by the optimum pH. There have been, so far reported that the presence of a ceramidase possessing the optimum pH in an acidic range has been found in mammalian tissues such as rat brain [*Biochemistry*, **8**, 1692-1698 (1969)], guinea pig epithelial cells [*J. Biol. Chem.*, **270**, 12677-12684 (1995)], human kidney [*Biochim. Biophys. Acta*, **398**, 125-131 (1975)], spleen [*Biochim. Biophys. Acta*, **1004**, 245-251 (1989)], fibroblasts [*Biochem. J.*, **205**, 419-425 (1982)], and epithelium [*FEBS Lett.*, **268**, 110-112 (1990)]; and human urine [*J. Biol. Chem.*, **270**, 11098-11102 (1995)], and the like.

In addition, it has been clarified that a bacterium belonging to the genus *Pseudomonas* produces a ceramidase, and this ceramidase is a ceramidase possessing optimum pH in an alkaline region [*J. Biol. Chem.*, **273**, 14368-14373 (1998)].

Among these ceramidases, amino acid sequences of the acidic ceramidase purified from human urine and nucleotide sequences of a gene encoding the ceramidase have been determined [*J. Biol. Chem.*, **271**, 33110-33115 (1996)]. In addition, an acidic ceramidase gene of a mouse has been obtained by utilizing its homology with the above-mentioned acidic ceramidase gene derived from human urine [*Genomics*, **50**, 267-274 (1998)].

However, since all of the ceramidase genes derived from mammals which have been reported encode acidic ceramidases, amino acid sequences and genomic structures of the neutral/alkaline ceramidase in mammals have been

completely unknown, so that biological functions of the neutral/alkaline ceramidase in higher organisms have not yet been elucidated at present.

In the studies on the elucidation of the *in vivo* functions of a ceramide, metabolic control therefor, diagnosis or treatment of a disease associated with the ceramide or the like, it is necessary to obtain detailed information concerning various enzymes associated with the ceramide, and the enzyme gene. However, as mentioned above, findings on the amino acid sequence and genes thereof of the neutral/alkaline ceramidase in mammals have not yet been obtained at present. Therefore, in order to develop the technique as described above pertaining to a ceramide, it is necessary to obtain some findings associated with a neutral/alkaline ceramidase, especially a gene thereof.

As mentioned above, although several reports have been made on cloning of ceramidase genes of a mammal, all of these reports are concerned with ceramidases possessing an activity in an acidic region, which cannot be expected to possess a homology with a ceramidase possessing an activity in a neutral/alkaline region. Therefore, it has been difficult to obtain a neutral/alkaline ceramidase gene as a homolog of a nucleotide sequence of the acidic ceramidase gene.

#### DISCLOSURE OF INVENTION

The present invention has been accomplished in view of the above-described prior art, and a first object of the present invention is to provide a neutral/alkaline ceramidase gene of a mammal. A second object of the present invention is to provide a method for producing a neutral/alkaline ceramidase gene in a high purity by a genetic engineering means, comprising using a

transformant resulting from incorporation of an expression vector carrying the gene. A third object of the present invention is to provide a polypeptide encoding the above-mentioned gene. A fourth object of the present invention is to provide an antisense DNA and an antisense RNA which are complementary to the gene of the present invention or a part thereof. A fifth object of the present invention is to provide a synthetic oligonucleotide probe or primer capable of specifically hybridizing to the gene of the present invention. A sixth object of the present invention is to provide an antibody or a fragment thereof specifically binding to the polypeptide. A seventh object of the present invention is to provide a method for detecting the above-mentioned ceramidase or a gene thereof, and a kit used therefor. An eighth object of the present invention is to provide a method of controlling an amount of ceramide in the cell or in the tissue.

The present inventors have succeeded in isolating a neutral/alkaline ceramidase from liver of a mouse, a mammalian, and isolating the genes. In addition, they have succeeded in elucidating the structure of the neutral/alkaline ceramidase of a mammal including human by using techniques such as hybridization or polymerase chain reaction (PCR). Further, they have also succeeded in conveniently producing a neutral/alkaline ceramidase in a high purity by genetic engineering techniques by using the genes. Thus, the present invention has been perfected thereby.

Specifically, the gist of the present invention relates to:

[1] a gene having a nucleotide sequence of a nucleic acid selected from the group consisting of:

(A) a nucleic acid encoding a polypeptide having the amino acid sequence of SEQ ID NO: 14 of the Sequence Listing or a partial sequence thereof, the

- polypeptide possessing a ceramidase activity;
- (B) a nucleic acid having a nucleotide sequence of SEQ ID NO: 15 of the Sequence Listing or a partial sequence thereof and encoding a polypeptide possessing a ceramidase activity;
- 5 (C) a nucleic acid encoding a polypeptide consisting of an amino acid sequence resulting from deletion, addition, insertion or substitution of at least one amino acid residue in the amino acid sequence of SEQ ID NO: 14 of the Sequence Listing, the polypeptide possessing a ceramidase activity;
- 10 (D) a nucleic acid consisting of a nucleotide sequence resulting from deletion, addition, insertion or substitution of at least one base in the nucleotide sequence of SEQ ID NO: 15 of the Sequence Listing and encoding a polypeptide possessing a ceramidase activity;
- (E) a nucleic acid capable of hybridizing to a complementary strand of a  
15 nucleic acid of any one of the above (A) to (D), under stringent conditions, and encoding a polypeptide possessing a ceramidase activity; and
- (F) a nucleic acid having a nucleotide sequence different from the nucleic acid of any one of above (A) to (E) via degeneracy and encoding a polypeptide possessing a ceramidase activity;
- 20 [2] a recombinant DNA comprising the gene of item [1] above;
- [3] an expression vector for a microorganism, an animal cell or a plant cell, comprising the gene of item [1] above or the recombinant DNA of item [2] above;
- [4] a transformant carrying the expression vector of item [3] above;
- 25 [5] a method for producing a polypeptide possessing a ceramidase activity,

characterized by culturing the transformant of item [4] above under conditions appropriate for expression of the ceramidase gene and production of the polypeptide encoded by the gene, and collecting a polypeptide possessing a ceramidase activity from the resulting culture;

5 [6] a polypeptide having the amino acid sequence of SEQ ID NO: 14 of the Sequence Listing or a partial sequence thereof and possessing a ceramidase activity;

[7] a polypeptide possessing a ceramidase activity, encoded by the gene of item [1] above;

10 [8] an antisense DNA which is complementary to the gene of item [1] above or a part thereof;

[9] an antisense RNA which is complementary to the gene of item [1] above or a part thereof;

[10] an expression vector comprising the antisense DNA of item [8] above;

15 [11] an oligonucleotide probe or primer, capable of specifically hybridizing to the gene of item [1] above or a complementary strand thereof;

[12] an antibody or a fragment thereof, capable of specifically binding to the polypeptide of item [6] or item [7] above;

20 [13] a method for detecting a gene encoding a polypeptide possessing a ceramidase activity, comprising using the oligonucleotide probe or primer of item [11] above;

[14] a kit for the use in detection of a gene encoding a polypeptide possessing a ceramidase activity, comprising the oligonucleotide probe and/or primer of item [11] above;

25 [15] a method for detecting a polypeptide possessing a ceramidase activity,

comprising using the antibody or a fragment thereof of item [12] above;  
[16] a kit for the use in detection of a polypeptide possessing a ceramidase  
activity, comprising the antibody or a fragment thereof of item [12] above; and  
[17] a method of controlling an amount of ceramide in a cell and/or in a tissue,  
5 characterized by introducing the gene of item [1] above or an antisense nucleic  
acid thereof into the cell and/or the tissue, thereby controlling the amount of  
ceramide in the cell and/or in the tissue.

#### BRIEF DESCRIPTION OF THE DRAWINGS

10 Figure 1 is a graph showing optimum pH of the ceramidase.

Figure 2 is a restriction endonuclease map of a DNA fragment comprising  
the ceramidase gene.

#### BEST MODE FOR CARRYING OUT THE INVENTION

15 Since information such as consensus sequences for a neutral/alkaline  
ceramidase has not been clarified, the present inventors have obtained the amino  
acid information by isolating the above-mentioned neutral/alkaline ceramidase,  
whereby a gene encoding the above-mentioned ceramidase could be isolated for  
the first time.

20 From the above findings, there could be obtained a surprising finding that  
the amino acid sequence of the neutral/alkaline ceramidase derived from a mouse  
liver of the present invention has a low homology with the amino acid sequence  
of a known alkaline ceramidase [previously mentioned, *J. Biol. Chem.*, **273**,  
14368-14373 (1998)] derived from a bacterium belonging to the genus  
25 *Pseudomonas* (*Pseudomonas aeruginosa*).

The ceramidase of the present invention has been clarified for the first time as a neutral/alkaline ceramidase derived from a mammal. Therefore, the present invention is even more useful in the developments of elucidation of the *in vivo* functions of a ceramide, metabolic control therefor, and diagnosis and treatment of diseases associated with a ceramide, and the like, as compared to the known alkaline ceramidase derived from the known bacterium of the genus *Pseudomonas*.

The present invention will be described hereinbelow.

(1) Polypeptide Possessing Ceramidase Activity

In the present specification, the phrase "polypeptide possessing a ceramidase activity" (which may be simply referred to as "ceramidase" in the present specification) refers to an enzyme possessing an activity of hydrolyzing a ceramide to generate a sphingoid and a fatty acid as mentioned above. In addition, the term "neutral/alkaline ceramidase" refers to a ceramidase possessing an optimum pH at a pH higher than the acidic range.

As one example thereof, enzymologically chemical and physicochemical characteristics of the isolated and purified, neutral/alkaline ceramidase derived from a mouse liver in the present invention will be described.

1. Action

The ceramidase of the present invention acts to hydrolyze a ceramide, thereby generating a sphingoid and a fatty acid.

The activity of the ceramidase can be determined in accordance with the method described, for instance, in *J. Biol. Chem.*, **275**, 3462-3468 (2000).

Concretely, a reaction mixture prepared by dissolving 550 pmol of C12-NBD-ceramide [*Anal. Biochem.*, **263**, 183-188 (1998)], 1.0% (W/V) sodium cholate and a suitable amount of an enzyme (ceramidase) in 20  $\mu$ l of 25 mM Tris-hydrochloric acid buffer (pH 7.5) is incubated at 37°C for 30 minutes. The reaction mixture is incubated in a boiling water bath for 5 minutes, thereby stopping the reaction. The resulting reaction mixture is further evaporated to dryness under reduced pressure. The dried solid is dissolved in chloroform/methanol = 2/1 (V/V), and developed by silica gel thin layer chromatography (developing solvent: chloroform/methanol/25% aqueous ammonia = 90/20/0.5 (V/V/V)). Thereafter, the C12-NBD-fatty acid generated by the above-mentioned reaction is quantified by using CS-9300 Chromatoscanner (manufactured by Shimadzu Corporation) at an excitation wavelength of 475 nm and a fluorescent wavelength 525 nm. One unit (U) of this enzyme (ceramidase) is defined as an amount of the enzyme required for releasing 1 micromol of the C12-NBD-fatty acid, per one minute under the above-mentioned conditions, resulting from hydrolysis of the C12-NBD-ceramide.

## 2. Substrate Specificity

Five milliunits of the ceramidase of the present invention is acted on 100 pmol of various kinds of sphingolipids, in which the fatty acid moiety is labeled with  $^{14}$ C-radioactive isotope, in 20 ml of 25 mM Tris-hydrochloric acid buffer, pH 7.5, containing 1% sodium cholate at 37°C for 24 hours. The reaction mixture is developed on silica gel thin layer chromatography. Thereafter, the  $^{14}$ C-labeled sphingolipids and the  $^{14}$ C-labeled fatty acids generated by the enzymatic reaction are detected and quantified by Imaging Analyzer BAS1000

(manufactured by Fuji Photo Film). The degradation ratio is calculated from the obtained values. The substrate specificities of the ceramidase of the present invention are shown in Table 1.

As shown in Table 1, the ceramidase of the present invention shows substrate specificities of: (1) hydrolyzing various N-acylsphingosines; (2) not acting on galactosylceramide, sulfatide, GM1a, or sphingomyelin; (3) acting favorably on a ceramide containing sphingenin (d18:1) than a ceramide containing sphinganine (d18:0); (4) being less likely to act a ceramide containing phytosphingosine (t18:0); and the like.

Table 1

| Substrate (Structure)                                  | Degradation Ratio (%) |
|--|-----------------------|
| N-Lauroylsphingosine (C12:0/d18:1)                     | 63                    |
| N-Palmitoylsphingosine (C16:0/d18:1)                   | 93                    |
| N-Stearoylsphingosine (C18:0/d18:1)                    | 83                    |
| N-Palmitoylsphinganine (C16:0/d18:0)                   | 59                    |
| N-Stearoylsphinganine (C18:0/d18:0)                    | 40                    |
| N-Palmitoylphytosphingosine (C16:0/t18:0)              | 5                     |
| N-Stearoylphytosphingosine (C18:0/t18:0)               | 2                     |
| NBD-N-Dodecanoylsphingosine (NBD-C12:0/d18:1)          | 97                    |
| NBD-N-Hexanoylsphingosine (NBD-C6:0/d18:1)             | 2                     |
| Galactosylceramide (Galb1-1' Cer)                      | 0                     |
| Sulfatide (HS03-3Galb1-1' Cer)                         | 0                     |
| GM1a (Galb1-3GalNAcb1-4(NeuAca2-3)Galb1-4G1cb1-1' Cer) | 0                     |
| Sphingomyelin (Choline phosphate Cer)                  | 0                     |

### 3. Optimum pH

Sixteen milliunits of the ceramidase of the present invention is acted on 100 pmol of C12-NBD-ceramide in 20 ml of 3,3-dimethylglutaric acid, 50 mM Tris-(hydroxymethyl)aminomethane, 50 mM 2-amino-2-methyl-1,3-propanediol at 37°C for 24 hours. The resulting reaction mixture is developed by silica gel thin layer chromatography. Next, the NBD-labeled ceramide and the NBD-labeled fatty acid generated by the enzymatic reaction are detected and quantified at a detection wavelength of 525 nm by using the CS-9300 Chromatoscanner. The degradation ratio is calculated from the obtained values. Figure 1 is a graph showing the relationship between the degradation activity of the C12-NBD-labeled ceramide and pH, wherein the ordinate axis shows the degradation ratio (%), and the abscissa axis shows a reaction pH. As shown in the results of Figure 1, the optimum pH of the present ceramidase is from 7.0 to 8.0.

### 4. Temperature Stability

The ceramidase of the present invention does not show the reduced activity when treated in 20 mM Tris-hydrochloric acid (pH 7.5) buffer containing 0.1% polidocanol [trade name: Lubrol PX] at 37°C for 24 hours, but shows the reduced activity by a treatment at 60°C for 1 hour, to about 30% of the activity before the treatment.

### 5. Molecular Weight

The molecular weight of the ceramidase of the present invention is about 94 kDa on SDS-PAGE (under reducing conditions). In addition, the present enzyme digested by glycopeptidase F is about 73 kDa on SDS-PAGE (under

reducing conditions).

In the present specification, one example of the "polypeptide possessing a ceramidase activity" includes a polypeptide of a naturally occurring ceramidase derived from a mouse, having the amino acid sequence of SEQ ID NO: 14 of the Sequence Listing. Further, not only the polypeptide having the amino acid sequence, but also a polypeptide having an amino acid sequence with a mutation such as substitution, deletion, addition or insertion of one or more amino acids introduced into the amino acid sequence of SEQ ID NO: 14 of the Sequence Listing are encompassed in the term "polypeptide possessing a ceramidase activity," as long as the polypeptide can be found to possess a similar ceramidase activity in accordance with the activity determination by the method described above. In addition, in the above-mentioned mutation, two or more kinds of mutations may be introduced, as long as they are mutations by which the resulting polypeptide can exhibit a ceramidase activity. In the present specification, the term "amino acid sequence with a mutation introduced" encompasses any of amino acid sequences into which a mutation is artificially introduced and amino acid sequences having a naturally occurring mutation.

The ceramidase having the mutation can be concretely obtained, for instance, by selecting a polypeptide possessing a ceramidase activity for the mutated gene having a mutation in the nucleotide sequence (SEQ ID NO: 15) of the ceramidase gene described below, by the following steps:

- (a) incubating a gene expression product in a reaction mixture [composition: 550 pmol of C12-NBD-ceramide and 1.0% (W/V) sodium cholate in 20  $\mu$ l of 25 mM Tris-hydrochloric acid buffer (pH 6 to 9, preferably 6.5 to 8.5,

more preferably 7 to 8, especially preferably 7.5)] at 37°C for 30 minutes, to react the mixture; and

(b) detecting generation of a C12-NBD-fatty acid in the reaction product.

5 (2) Ceramidase Gene

The ceramidase gene in the present invention refers to a gene having a nucleotide sequence of a nucleic acid encoding the above-mentioned  
“polypeptide possessing a ceramidase activity,” or a nucleic acid comprising a nucleotide sequence of the gene. One example thereof includes a gene having a  
10 nucleotide sequence of a nucleic acid encoding a polypeptide consisting of the amino acid sequence of SEQ ID NO: 14 of the Sequence Listing or a partial sequence thereof; and a gene having the nucleotide sequence of a nucleic acid consisting of a nucleotide sequence of SEQ ID NO: 15 of the Sequence Listing, or a partial sequence thereof, and the present invention is not limited to these  
15 exemplified. As described above, even a gene having a nucleotide sequence of a nucleic acid encoding a polypeptide consisting of a partial sequence of the amino acid sequence of SEQ ID NO: 14, or a gene having a nucleotide sequence of a nucleic acid consisting of a partial sequence of the nucleotide sequence of  
20 SEQ ID NO: 15 are encompassed in the scope of the present invention, as long as the gene encodes a polypeptide possessing a ceramidase activity. These genes are neutral/alkaline ceramidase genes derived from mouse liver, and the origin of the ceramidase gene of the present invention is not particularly limited, as long as the ceramidase activity of the gene product can be detected in the same manner as in the steps (a) and (b) described in item (1) above. The origin  
25 includes, for instance, mice, rats, humans, hamsters, guinea pigs, and the like.

In the present specification, the phrase "polypeptide consisting of a partial sequence (of the amino acid sequence)" means those in which the ceramidase activity can be detected by the steps (a) and (b) described in item (1) above.

Further, a gene encoding a ceramidase having a mutation, which can exhibit a similar ceramidase activity, is also encompassed in the present invention. For instance, even a gene having a nucleotide sequence of a nucleic acid encoding an amino acid sequence with a mutation such as deletion, addition, insertion or substitution of one or more amino acid residues introduced into the amino acid sequence of SEQ ID NO: 15 of the Sequence Listing is encompassed in the gene of the present invention, as long as the polypeptide encoded by the gene possesses a ceramidase activity. Even a naturally occurring gene having a mutation as well as an artificially prepared gene described above are encompassed in the scope of the present invention, as long as the gene is a gene having a nucleotide sequence of a nucleic acid encoding a polypeptide possessing an activity for a neutral/alkaline ceramidase.

As the method for preparing a gene into which a mutation is artificially introduced as described above, for instance, the following method is used.

As a method of introducing a random mutation, there can be employed, for instance, a method for generating a transition mutation in which cytosine base is substituted by uracil base with a chemical treatment using sodium hydrogen sulfite [*Proceedings of the National Academy of Sciences of the USA*, 79, 1408-1412 (1982)]; a method for lowering an accuracy of incorporation of nucleotide during DNA synthesis by carrying out PCR in a reaction mixture containing manganese [*Anal. Biochem.*, 224, 347-353 (1995)]; and the like.

As a method for introducing site-directed mutagenesis, there can be

employed, for instance, a method utilizing amber mutation [gapped duplex method, *Nucleic Acids Research*, 12, 9441-9456 (1984)]; a method of utilizing a *dut* (dUTPase) gene and *ung* (uracil-DNA glycosylase) gene deficient host [Kunkel method, *Proceedings of the National Academy of Sciences of the USA*, 82, 488-492 (1985)]; a method by PCR utilizing amber mutation (WO 98/02535); and the like. Various kinds of kits for introducing site-directed mutagenesis to a desired gene by these methods are commercially available, and a gene resulting from introduction of a desired mutation can be readily obtained by utilizing the kits.

In addition, the nucleic acid capable of hybridizing to a complementary strand of the above-mentioned nucleic acid under stringent conditions, the gene having a nucleotide sequence of a nucleic acid encoding a polypeptide possessing a ceramidase activity is also encompassed in the gene of the present invention. The ceramidase activity can be detected, for instance, by the steps (a) and (b) described in item (1) above.

Here, the term "stringent conditions" refers to, for instance, the following conditions. Concretely, the term refers to conditions of carrying out incubation at 50°C for 4 hours to overnight in 6 x SSC (wherein 1 x SSC is 0.15 M NaCl, 0.015 M sodium citrate, pH 7.0) containing 0.5% SDS, 5 x Denhardt's [0.1% bovine serum albumin (BSA), 0.1% polyvinyl pyrrolidone, 0.1% Ficoll 400], and 100 µg/ml salmon sperm DNA.

Also, the details of the hybridization manipulations are described, for instance, in *Molecular Cloning: A Laboratory Manual*, 2nd Ed., published by Cold Spring Harbor Laboratory in 1989, edited by T. Maniatis et al.

The gene having a nucleotide sequence different from the nucleic acid

mentioned above via a degenerated genetic code is also encompassed in the ceramidase gene of the present invention.

The polypeptide encoded by the gene of the present invention is encompassed in the present invention, as long as its ceramidase activity can be detected by the steps (a) and (b) described in item (1) above.

According to the gene of the present invention, there is further provided a recombinant DNA suitable for various purposes used. Here, the term "recombinant DNA" refers to a DNA carrying the gene of the present invention, obtained by genetic engineering technique.

The recombinant DNA carrying the ceramidase gene of the present invention is ligated to a known vector or the like, whereby an expression vector inserted with the ceramidase gene in an expressible state can be prepared. Such an expression vector is also encompassed in the present invention.

In the present invention, an expression vector refers to a vector which is constructed such that the gene or recombinant DNA mentioned above is inserted and expressed in desired host cells. In addition, a vector in which an antisense DNA as described below is inserted is also encompassed in the expression vector of the present invention. The vector to be inserted includes plasmid vectors, phage vectors, viral vectors, and the like. As the plasmid vector, commercially available products such as pUC18, pUC19, pBluescript, and pET can be suitably used, and as the phage vectors, commercially available products of lambda phage vectors such as  $\lambda$ gt10 and  $\lambda$ gt11 can be suitably used, without being limited thereto. As the viral vectors, retroviral vector, adenoviral vector, vaccinia viral vector, adeno-associated viral vector, and the like can be used, without being limited thereto. These vectors are appropriately selected in accordance with the

host cells used. Each of these vectors may appropriately carry a factor such as an inducible promoter, a selectable marker gene, or a terminator.

In addition, in order to facilitate isolation and purification, a vector carrying a sequence capable of expressing as His tag or GST fusion protein may be used depending upon its use. In this case, GST (glutathione S-transferase) fusion protein vector carrying an appropriate promoter (for instance, *lac*, *tac*, *trc*, *trp*, CMV, SV40 early promoter, or the like) which functions in a host cell (for instance, pGEX4T), a vector carrying a tag (for instance, Myc, HisA, or the like) sequence or the like can be used.

### (3) Transformant Harboring Ceramidase Gene

The transformant of the present invention, namely the cells capable of expressing the ceramidase gene of the present invention, can be obtained by transforming a host with an expression vector in which the ceramidase gene of the present invention is inserted. The host used can be appropriately selected depending upon the purposes used of the desired ceramidase, and microorganisms such as *Escherichia coli*, yeasts, animal cells, plant cells, animal individuals, plant individuals, and the like. Concretely, *Escherichia coli* includes HB101 strain, C600 strain, JM109 strain, DH5 $\alpha$  strain, DH10B strain, XL-1BlueMRF' strain, TOP10F strain and the like of the *Escherichia coli* K-12 derivative. Also, the yeast cells include *Saccharomyces cerevisiae* and the like. The animal cells include L, 3T3, FM3A, CHO, COS, Vero, Hela, and the like. The plant cells include tobacco BY2 and the like.

As a method for transducing an expression vector into a host, a method, including, for instance, described in *Molecular Cloning: A Laboratory Manual*,

2nd Ed., 249-254 may be employed. Next, in order to select a transformant expressing a desired gene, the characteristics of the expression vector are utilized. For instance, in a case where a plasmid vector is pBluescript and a host cell *Escherichia coli*, a colony having ampicillin resistance on a plate containing ampicillin is selected, or a colony having ampicillin resistance and showing white color on a plate containing ampicillin, 5-bromo-4-chloro-3-indolyl- $\beta$ -D-galactoside (X-Gal) and isopropyl- $\beta$ -D-thiogalactopyranoside (IPTG) is selected, thereby selecting a colony in which a foreign gene is transduced.

The polypeptide possessing a ceramidase activity can be produced by culturing the transformant of the present invention under generally employed conditions. In some cases, the codon usage differs depending upon a host in which the gene of the present invention is expressed, so that expression is suppressed. In this case, the codon used in the gene of the present invention may be changed to a codon suitable to each host. In addition, the above-mentioned expression vector is not limited only to those vectors derived from plasmids, and vectors derived from phages, cosmids, and the like may also be used, as long as the object of the present invention is not hindered. A vector capable of inducing and expressing a foreign gene, a vector capable of expressing as a fusion protein with a reporter gene product, and the like are desirable, from the viewpoint of readily and massively producing the polypeptide of the present invention.

The expression of the ceramidase can be confirmed by determining the ceramidase activity. The activity can be determined by, for instance, the method described in *J. Biol. Chem.*, **275**, 3462-3468 (2000) using a cell extract of the transformant as a sample, for instance, similar procedures to the steps (a) and (b) described in item (1) above. In addition, the expression of the ceramidase can be

confirmed by determining the amount of a ceramide in the cell. The above-mentioned amount of ceramide can be determined, for instance, by the method described in *Analytical Biochemistry*, 244, 291-300 (1997). Also, an antibody against the ceramidase can be used. In a case where the ceramidase is expressed as a fusion body with another polypeptide (the polypeptide excluding the ceramidase of the present invention), an antibody against this polypeptide moiety, the polypeptide excluding the ceramidase of the present invention, may be used. In a case where an antibody is used, the ceramidase can be detected by, for instance, subjecting a cell extract of the transformant to electrophoresis on an SDS-polyacrylamide gel, and thereafter transferring the electrophoresed gel on a polyvinylidene fluoride (PVDF) membrane, and detecting with the antibody on this membrane.

#### (4) Method for Producing Polypeptide Possessing Ceramidase Activity

The present invention also provides a method for producing a polypeptide possessing a ceramidase activity, comprising the steps of culturing the above-mentioned transformant under conditions appropriate for expression of the ceramidase gene of the present invention and production of the polypeptide encoding the gene, and collecting a polypeptide possessing a ceramidase activity from the resulting culture. The method for culturing the transformant is not particularly limited, and an appropriate one can be selected from a known culturing method appropriate for the host used.

In the production method of the present invention, in a case where the above-mentioned transformant is a microorganism or a cultured cell, the ceramidase can be efficiently produced by determining optimum conditions for

expression of the ceramidase in a medium composition, a pH of a medium, a culturing temperature, and a culturing time, as well as the amount of an inducer used, and the using time period, and the like.

5 A general method is employed in the purification of the ceramidase from the culture of the transformant. In a case where the transformant intracellularly accumulates the ceramidase as in the case of *Escherichia coli*, after the termination of culture, the transformant cells are harvested by centrifugation, and the resulting cells are disrupted by sonication or the like, and thereafter centrifuged or the like to give a cell-free extract. The cell-free extract used as a  
10 starting material can be purified by salting-out as well as a general protein purification method such as various kinds of chromatographies such as ion-exchange chromatography, gel filtration chromatography, hydrophobic chromatography, and affinity chromatography. The expression product may be extracellularly secreted depending upon the host-vector system used in some  
15 cases. In such a case, the purification can be similarly carried out from culture supernatant.

According to the method of the present invention, in a case where the ceramidase is produced intracellularly, a desired ceramidase may coexist intracellularly with impurities such as various kinds of enzymes and proteins.  
20 Since the impurities are present in a trace amount as compared to the amount of the expressed ceramidase, there is an excellent advantage that its purification is extremely facilitated. In addition, in a case where an extracellular secretion-type vector is used as a vector, the ceramidase is extracellularly secreted, so that the medium component or the like coexists in the fraction comprising the ceramidase.  
25 However, since the fraction usually contains almost no protein component which

would hinder the ceramidase purification, there is an excellent advantage in that, for instance, complicated separation and purification procedures which have been required in the purification of the ceramidase from a mouse liver are not necessitated.

5           In addition, in a case of a ceramidase derived from a fungus, there is a possibility that the enzyme itself has a sugar chain. A polypeptide possessing a ceramidase activity and having no sugar chain can be produced by using as a host cell a cell which does not possess a sugar chain-synthesizing ability, for instance, a mutant cell which has lost a sugar chain-synthesizing ability of a  
10   prokaryote such as *Escherichia coli*, *Bacillus subtilis* or *Actinomyces*, or an yeast, a fungus, an animal cell, an insect cell and a plant cell. Further, an enzyme having a sugar chain can be produced. In this case, a polypeptide possessing a ceramidase activity and having a sugar chain can be produced by using as a host cell a cell which possesses a sugar chain-synthesizing ability, for  
15   instance, an yeast, a fungus, an animal cell, an insect cell and a plant cell.

          Also, the expression product may form an insoluble inclusion body depending upon the host-vector system used. In this case, after the termination of culture, the transformant cells are harvested by centrifugation, and the resulting cells are disrupted by sonication or the like, and thereafter centrifuged  
20   or the like, thereby harvesting an insoluble fraction containing the inclusion body. After washing the inclusion body, the inclusion body is solubilized with an generally used protein solubilizing agent, for instance, urea, guanidine hydrochloride, or the like, and purified by various kinds of chromatographies such as ion-exchange chromatography, gel filtration chromatography,  
25   hydrophobic chromatography, and affinity chromatography as occasion demands.

Thereafter, refolding procedures employing dialysis or dilution method are carried out, whereby a preparation comprising a polypeptide possessing a ceramidase activity can be obtained. If this preparation is further purified by various kinds of chromatographies as occasion demands, a polypeptide  
5 possessing a ceramidase activity in a high purity can be obtained.

(5) Probe for Hybridization and Primer for PCR

The oligonucleotide probe or primer of the present invention is capable of specifically hybridizing to the gene of the present invention, or a complementary  
10 strand thereof. The oligonucleotide probe or primer is designed on the basis of the nucleotide sequence of the ceramidase gene of the present invention. For instance, the oligonucleotide probe or primer can be prepared by chemical synthesis by a general method. The nucleotide sequence for the oligonucleotide probe is not particularly limited. The nucleotide sequence is those capable of  
15 hybridizing to the above-mentioned ceramidase gene, or a nucleic acid having a nucleotide sequence complementary to the gene under stringent conditions. The above-mentioned term "stringent conditions" is not particularly limited. For instance, the term "stringent conditions" refers to conditions of incubating overnight at a temperature of  $[(T_m \text{ of the above-mentioned probe}) - 25^\circ\text{C}]$  in a  
20 solution containing 6 x SSC, 0.5% SDS, 5 x Denhardt's, and 100 mg/ml salmon sperm DNA, and the like. Also, the nucleotide sequence of the above-mentioned primer is not particularly limited, as long as the primer is annealed to the above-mentioned ceramidase gene or the gene having a nucleotide sequence complementary to the gene under usual reaction conditions for PCR so that the  
25 extension reaction by the DNA polymerase can be initiated.

*T<sub>m</sub>* of the oligonucleotide probe or primer can be calculated, for instance, by the following equation:

$$T_m = 81.5 - 16.6 (\log_{10}[\text{Na}^+]) + 0.41 (\%G + C) - (600/N)$$

wherein N is a chain length of the oligonucleotide probe or primer; and

%G + C is a content of guanine and cytosine residues in the oligonucleotide probe or primer.

In addition, when the chain length of the oligonucleotide probe or primer is shorter than 18 bases, *T<sub>m</sub>* can be deduced from a product of the contents of A + T (adenine + thymine) residues multiplied by 2°C, with a sum of a product of the contents of G + C residues multiplied by 4°C [(A + T) x 2 + (G + C) x 4].

The chain length of the above-mentioned oligonucleotide probe is not particularly limited. It is preferable that the chain length is 15 bases or more, more preferably 18 bases or more, from the viewpoint of preventing nonspecific hybridization.

In addition, as the primer of the present invention, there can be cited the nucleic acids having the same nucleotide sequences as those for the above-mentioned oligonucleotide probe. For instance, the primer can be prepared by, for instance, designing on the basis of the nucleotide sequence of the gene of the present invention, and chemically synthesizing it, and the like. The chain length of the primer is not particularly limited. For instance, the primer having a chain length of 15 to 40 bases can be used, especially one having a chain length of 17 to 30 bases can be suitably used. The above-mentioned primer can be used for various gene amplification methods such as PCR method, whereby the ceramidase gene of the present invention can be detected.

Also, as the above-mentioned oligonucleotide probe or primer, there may be used a nucleic acid obtained by fragmenting a nucleic acid encoding a naturally occurring ceramidase by an enzymatic treatment such as endonuclease treatment or exonuclease treatment, a physical treatment such as sonication, or the like, and subjecting the resulting fragment to separation and purification by various kinds of nucleic acid separation methods represented by agarose gel or the like. It is desired that the nucleic acid obtained as described above is derived from a region having a sequence characteristic of the ceramidase.

Further, in order to more readily detect the nucleic acid to be detected, the above-mentioned oligonucleotide probe or primer can be subjected to appropriate labeling in accordance with a known method to be used in the detection of the ceramidase gene of the present invention. The labeling is not particularly limited. The oligonucleotide probe or primer may be labeled with radioisotopes as well as various labels represented by fluorescent substances, and ligands such as biotin and digoxigenin.

A DNA having high homology to the ceramidase gene of the present invention can be cloned by screening a genomic DNA or cDNA derived from an organ other than a mouse liver or from an organism other than a mouse, or a genomic DNA library or cDNA library, by using the probe for hybridization of the present invention.

In addition, a DNA fragment having high homology to the ceramidase gene of the present invention can be detected by PCR method from a genomic DNA or cDNA derived from an organ other than mouse liver or derived from an organism other than mouse, or a genomic DNA library or cDNA library, by using the primer of the present invention, and further its full length gene can also

be obtained.

(6) Method for Detecting Gene

One of the big features of the method for detecting a gene of the present invention resides in that the gene in the sample to be detected is detected by using the above-mentioned oligonucleotide probe and/or primer.

In the method for detection of the present invention, the gene may be detected by hybridization method or the like by using the above-mentioned oligonucleotide probe, or the gene may be detected by DNA amplification method such as PCR method by using the above-mentioned primer.

In a case of hybridization using the oligonucleotide probe, the samples to be detected include, for instance, samples such as colonies and cultured cells of microorganisms, and tissue fragments, those obtained by immobilizing DNA or RNA in these samples onto a membrane, DNA or RNA extracted from these samples, and the like.

The hybridization can be carried out in accordance with a known method described in *Molecular Cloning: A Laboratory Manual*, 2nd Ed. and the like. The conditions for the hybridization can be appropriately determined by the  $T_m$  value of the probe used, the GC content of the target DNA, and the like. For instance, the conditions described in *Molecular Cloning: A Laboratory Manual*, 2nd Ed. mentioned above or the like can be applied.

In a case where the gene is detected by using the primer, the samples to be detected include, for instance, microorganism samples such as culture of microorganisms, colonies of microorganisms and bacterial cells of microorganisms; samples derived from a living body such as cultured cells,

tissues and tissue fragments; and the like. As these samples, for instance, isolated microorganisms and cultured cells may be used in the original states, or after being subjected to an appropriate treatment. In addition, solid samples such as tissues can be used by preparing an exudate or suspension. Also, supernatant of these samples, or those samples prepared by subjecting these samples to a cytolytic treatment such as a treatment with a surfactant, or supernatant thereof can be used. Further, the sample may be subjected to a procedure of removing other components in the sample within a range so as not to impair the nucleic acid to be detected.

10 In a case where the detection is carried out by PCR method by using the above-mentioned primer, the PCR conditions can be appropriately selected in accordance with the  $T_m$  value of the primer used, the length of the region to be amplified and detected, and the like. In PCR, the desired gene can be detected by confirming the presence or absence of the amplified product. The method for confirming the presence or absence of amplification is not particularly limited. For instance, the confirmation can be made by, for instance, subjecting a reaction mixture for nucleic acid amplification to agarose gel electrophoresis, staining the gel with an appropriate nucleic acid staining reagent, such as ethidium bromide or SYBER Green I, subjecting the gel to irradiation with ultraviolet rays, and detecting the presence or absence of the band. The detection of the band may be observed by naked eyes, or the detection can be made, for instance, by using a fluorescent image analyzer or the like.

25 In the method for detecting the gene of the present invention, in order to increase the detection sensitivity, the above-mentioned probe and primer may be used together. For instance, the gene can be detected in high sensitivity and

accurately by amplifying the ceramidase gene existing in a trace amount in the sample by PCR method with the above-mentioned primer, and thereafter hybridizing the gene with the probe.

In a case where the ceramidase gene is detected by the method for detection of the present invention and the amount of the gene is further determined, the amount of the gene can be determined by quantifying an intensity of the signal ascribed to the hybridized probe, a fluorescent intensity of the band ascribed to a product amplified with the primer, or the like. The expression amount of the desired gene can be examined by quantifying the amount by using mRNA as an object to be determined.

In addition, in the method for detection of the present invention, the detection can be carried out more conveniently by using the kit for the use in detection of the gene of the present invention. Such a kit is also encompassed in the present invention. One of the features of the above-mentioned kit resides in that the kit comprises the above-mentioned oligonucleotide probe and/or the above-mentioned primer. The above-mentioned kit may contain various components used in the detection procedures. For instance, in a case of a kit comprising an oligonucleotide probe, there may be contained various kinds of reagents for hybridization representatively exemplified by a membrane for immobilizing a nucleic acid, a hybridization buffer, and the like. Also, in a case of a kit comprising a primer, there may be contained reagents for PCR representatively exemplified by thermostable DNA polymerases, dNTP mixed solutions, buffers for PCR, and the like. Further, there may be contained reagents for detecting a probe or an amplified DNA, media for proliferating microorganisms, media for culturing cells, reagents for extracting nucleic acids

from a sample, and the like.

(7) Antibody or Fragment Thereof Specifically Binding to Polypeptide  
Possessing Ceramidase Activity

5 The antibody or a fragment thereof specifically binding to the polypeptide  
of the present invention is not particularly limited, as long as the antibody or a  
fragment thereof possesses an ability of specifically binding to the polypeptide.  
The antibody may be any of polyclonal antibodies and monoclonal antibodies.  
Further, antibodies modified by known techniques or antibody derivatives, for  
10 instance, humanized antibodies, Fab fragments, single-chain antibodies, and the  
like, can also be used. The antibody of the present invention can be readily  
prepared by appropriately immunizing a rabbit, a rat or a mouse using all or a  
part of the polypeptide of the present invention in accordance with the method  
described in, for instance, *Current Protocols in Immunology*, edited by John E.  
15 Coligan, published by John Wiley & Sons, Inc., 1992. The antibody thus  
obtained is purified and thereafter treated with a peptidase or the like, to give an  
antibody fragment. In addition, the antibody can be prepared by genetically  
engineering means. Further, the antibody or a fragment thereof of the present  
invention may be subjected to various modifications in order to facilitate the  
20 detection by enzyme immunoassay, fluoroimmunoassay, luminescent  
immunoassay, or the like.

The antibody or a fragment thereof mentioned above encompasses those  
which are capable of specifically binding to a certain partial fragment of the  
polypeptide.

25 The use of the resulting antibody or a fragment thereof includes

applications to detection of ceramidase-producing bacteria, detection of ceramidase-expressed cell lines, detection of ceramidase proteins in the cultured cell or in the tissue, affinity chromatography, screening of an expression product of various kinds of libraries (genomic DNA or cDNA), pharmaceuticals, diagnostic agents, reagents for researches, and the like.

(8) Method for Detecting Polypeptide

A feature of the method for detecting the polypeptide of the present invention resides in that the polypeptide possessing a ceramidase activity is detected by the antibody or a fragment thereof mentioned above.

In the present invention, as the samples to be detected, there can be used, for instance, cultures of microorganisms and animal cells, tissue fragments, cell disruptions of microorganisms and animal cells, extracts or washings of tissues such as skins, and protein samples such as membranes immobilized with proteins derived from microorganisms, animal cells and tissues.

As to the detection of the specific binding of the antibody or a fragment thereof to the above-mentioned polypeptide, a known method can be utilized, and such a method includes, for instance, enzyme immunoassay, fluoroimmunoassay, luminescent immunoassay, and the like.

In the method for detecting the polypeptide of the present invention, the detection can be more conveniently carried out by using the kit for the use in the detection of the polypeptide of the present invention. Such a kit is also encompassed in the present invention. A feature of the above-mentioned kit resides in that the kit comprises the above-mentioned antibody or a fragment thereof. In addition, the kit may contain a reaction buffer, a labeled secondary

antibody, a developing reagent, and the like.

(9) Antisense DNA and Antisense RNA

In the present invention, each of the terms "antisense DNA" and  
5 "antisense RNA" refers to those having a nucleotide sequence complementary to  
the ceramidase gene of the present invention or a part thereof, which suppresses  
or controls expression (transcription, translation) of the genetic information from  
the gene by forming a double strand with an endogenous ceramidase gene  
(genomic DNA and mRNA). The length of the antisense DNA or antisense  
10 RNA can be changed depending upon the specificity of the nucleotide sequence  
and the method for transducing the nucleotide sequence into a cell. The  
antisense DNA or antisense RNA can be prepared by artificially synthesizing  
with a synthesizer; expressing a gene in an opposite direction (direction of  
antisense) of the usual direction by an enzymatic reaction with the gene of the  
15 present invention as a template; or the like. In a case where expression of the  
antisense RNA is desired in a living body, an expression vector ligated to the  
gene of the present invention in an opposite direction of the usual direction is  
constructed, and the resulting expression vector may be transduced into a living  
body.

20 For instance, numerous antisense techniques such as suppression of  
proliferation of HIV utilizing *tat* gene [*Nucleic Acids Research*, **19**, 3359-3368  
(1991)] or *rev* gene [*Proceedings of the National Academy of Sciences of the  
USA*, **86**, 4244-4248 (1989)] have been known. Therefore, according to these  
25 methods, expression of the endogenous ceramidase gene can be suppressed or  
controlled by using the antisense DNA or antisense RNA of the present invention.

In addition, the antisense DNA or antisense RNA of the present invention can be utilized as a research reagent for *in situ* hybridization or the like.

(10) Control of Amount of Ceramide in Cell or in Tissue Using Ceramidase Gene or Antisense Nucleic Acid Thereof

By the gene of the present invention, a method of controlling an amount of a ceramide in a cell and/or in a tissue can be further provided. Such a method of control is also encompassed in the present invention. One of the features of the method of controlling an amount of a ceramide in a cell and/or in a tissue of the present invention resides in that the ceramidase gene of the present invention is introduced into the cell and/or into the tissue, thereby controlling the amount of a ceramide in the cell and/or in the tissue.

Concretely, in the method of control of the present invention, the ceramidase gene of the present invention is introduced into the cell and/or into the tissue, and the ceramide is degraded in a cell or in a tissue by an action of the ceramidase expressed by the gene. On the other hand, the ceramidase gene of the present invention is introduced into the cell or into the tissue so as to generate the antisense nucleic acid of the gene, for instance, an antisense RNA, whereby the ceramidase activity in the cell or in the tissue is lowered, so that the degradation of the ceramide can be suppressed. In addition, in a case where the amount of the ceramide is suppressed, the antisense nucleic acid of the ceramidase gene of the present invention, namely the antisense DNA or antisense RNA of the present invention, may be introduced in a cell or in a tissue in an intact form.

As the method for introducing the ceramidase gene or an antisense nucleic

acid thereof mentioned above in a cell or in a tissue, a known method can be used, and methods for physically introducing a gene such as electroporation method and particle gun method, or a method for introducing a gene using a viral vector can be utilized. The viral vector which can be used in the method of the present invention is not particularly limited. For instance, the gene can be introduced by using retroviral vector, adenoviral vector, vaccinia viral vector, adeno-associated viral vector, or the like.

According to the method of control of the present invention, the ceramidase gene or an antisense nucleic acid thereof of the present invention is introduced in a cell or in a tissue to control the amount of a ceramide in the cell and/or in the tissue, whereby exhibiting excellent effects such that a disease caused by an abnormal amount of a ceramide can be treated, and that a model animal suffering from a disease with abnormal ceramide metabolism can be prepared. The "disease caused by an abnormal amount of a ceramide" is not particularly limited, and includes, for instance, Farber's disease and the like.

The method for obtaining a ceramidase gene derived from a mouse liver will be explained hereinbelow.

1) First, a membrane fraction is prepared from a homogenate of a mouse liver, and suspended in a sucrose-EDTA solution, and the suspension is frozen and thawed, and thereafter centrifuged, to give supernatant (crude enzyme extract). A ceramidase preparation homogeneously purified can be obtained from the resulting crude enzyme extract by combining known protein purification methods, for instance, various kinds of chromatographies. As the chromatographies which can be used in the above-mentioned purification, anion

exchange chromatography, hydrophobic chromatography, chelating chromatography, gel filtration chromatography, and the like can be used.

2) Next, as the information for preparing a probe for cloning the ceramidase gene, the partial amino acid sequences of the ceramidase are examined. The N-terminal amino acid sequence of the ceramidase can be found by subjecting the above-mentioned purification ceramidase preparation itself to amino acid sequencing by Edman degradation method [*J. Biol. Chem.*, **256**, 7990-7997 (1981)]. In addition, the partial amino acid sequence of the internal ceramidase can be obtained by purifying an appropriate peptide fragment from a peptide mixture resulting from digestion of the purified enzyme preparation with a protease having a high substrate specificity, for instance, lysylendopeptidase, N-tosyl-L-phenylalanyl chloromethyl ketone (TPCK)-trypsin, and the like, and subjecting the resulting fragment to amino acid sequencing.

3) On the basis of the information of the amino acid sequence thus clarified, an oligonucleotide to be used for the probe for hybridization, or used for the primer for PCR is designed, to clone the ceramidase gene of the present invention. In order to carry the cloning, PCR or hybridization method generally employed is utilized. PCR method can be carried out in accordance with the method described in *PCR Technology*, edited by Erlich, H.A., published by Stockton Press, 1989. The hybridization method can be carried out, for instance, in accordance with a known method described in *Molecular Cloning: A Laboratory Manual*, 2nd Ed.

4) As to the DNA fragment obtained by the hybridization or PCR mentioned above, by decoding its nucleotide sequence, the amino acid sequence which can be encoded therein can be known. Whether or not the above-mentioned DNA

fragment is a fragment of the ceramidase gene can be confirmed by comparing the sequence with the partial amino acid sequence of the ceramidase obtained in item 2) above.

5) In a case where the DNA fragment obtained by hybridization or PCR in item 3) is a part of the ceramidase gene, a DNA fragment comprising a gene encoding a full length ceramidase can be obtained by repeating the procedures of item 3), or alternatively, preparing a new probe or primer on the basis of the nucleotide sequence of the DNA fragment obtained in item 3), and carrying out hybridization or PCR by using the probe or primer,.

6) An expression vector is constructed by ligating the gene encoding a full length ceramidase thus obtained with an appropriate vector, and then the transformant in which the expression vector is transduced is prepared. The transformant is cultured, and the ceramidase activity in the resulting culture is examined, whereby confirmation can be made such that the resulting gene is one encoding the ceramidase.

However, in the cloning of the ceramidase gene of the present invention, a probe DNA appropriate for screening of the library by hybridization method could not be designed on the basis of the partial amino acid sequence information obtained in the present invention. In addition, although various PCR primers were designed by designing from each of the partial amino acid sequences and the nucleotide sequences of vector for the use in the library preparation, and the resulting primers were used in various combination to carry out PCR, specific amplification was not found, the only amplification being found alone in the combination of two kinds of primers designed on the basis of the partial amino acid sequence C-53 (shown in the amino acid sequence of C-53

in SEQ ID NO: 3 of Sequence Listing). However, the amplified DNA fragment P-1 (shown in the nucleotide sequence of P-1 in SEQ ID NO: 6 of Sequence Listing) was as short as 68 bp, so that the amplified fragment itself could not be used as a probe for library screening by hybridization method. Therefore, a gene fragment of a size of 335 bp, which is thought to be appropriate for the first time as a probe for library screening by hybridization method, has been successfully obtained for the first time by further designing a primer for PCR on the basis of the sequence of P-1, and at the same time carrying out PCR in combination of the primer and a primer designed on the basis of the nucleotide sequence of a vector used in the preparation of the library.

Further, a gene encoding a full length ceramidase can be cloned by screening cDNA library derived from a mouse liver with the above-mentioned fragment of a size of 335 bp as a probe. Also, the genomic DNA of a ceramidase of the present invention can be obtained by screening a genomic DNA library derived from a mouse liver.

The entire nucleotide sequence of the produced ceramidase gene from a mouse liver thus obtained is shown in SEQ ID NO: 12 of Sequence Listing, and the amino acid sequence of the polypeptide encoded thereby is shown in SEQ ID NO: 13 of Sequence Listing. It has been found that the enzyme is processed to a mature enzyme in which the N-terminal part of the peptide of the gene is removed *in vivo* on the bases of this amino acid sequence and the N-terminal amino acid sequence of the gene. The amino acid sequence of this mature ceramidase and the nucleotide sequence encoded by the sequence are shown in SEQ ID NOs: 14 and 15 of Sequence Listing, respectively. Each of the above-mentioned amino acid sequence and nucleotide sequence does not have

homology with each of the known amino acid sequence and nucleotide sequence of a ceramidase derived from a mammal. In other words, the ceramidase gene provided by the present invention consists of a completely novel sequence, which is irrelevant to the known ceramidase gene.

5 As described above, according to the present invention, there are provided a primary structure and a genetic structure of the ceramidase derived from a mouse liver. Further, there can be carried out a method for producing a polypeptide possessing a ceramidase activity inexpensively and in a high purity by genetic engineering means.

10 In addition, the oligonucleotide probe or primer capable of specifically hybridizing to the ceramidase gene of the present invention is useful in searching, detection, amplification and the like of the ceramidase gene of the present invention. The antibody or a fragment thereof specifically binding to the polypeptide of the present invention is useful in detection, identification,  
15 purification, and the like of a ceramidase.

In addition, according to the method of controlling an amount of a ceramide in a cell and/or in a tissue of the present invention, the amount of a ceramide in the cell and/or in the tissue can be controlled by introducing the ceramidase gene of the present invention or its antisense nucleic acid into the cell  
20 and/or into the tissue. Therefore, such a method of control is useful in the treatment of a disease caused by an abnormal ceramide amount, for instance, but not being particularly limited thereto, a treatment of a disease such as Farber's disease.

25 The present invention will be concretely explained by the examples,

without by no means intending to limit the scope of the present invention to these examples.

#### Example 1 Purification of Ceramidase

5 In 300 ml of a 0.25 M sucrose solution containing 1 mM EDTA (sucrose-EDTA solution) was homogenized 181 g of livers excised from 105 Sea/ddY mice (produced by Seiwa Experimental Animals). The resulting homogenate was centrifuged at 600 x g for 10 minutes, and thereafter the supernatant was collected. The resulting supernatant was further centrifuged at 2700 x g for 10 30 minutes, and the precipitates were collected.

The precipitated fraction was suspended in 480 ml of the sucrose-EDTA solution to give a suspension. The resulting suspension was frozen at -80°C, and thereafter thawed under running water. This freezing-thawing treatment was repeated twice. Thereafter, the treated suspension was centrifuged at 105000 x g 15 for 90 minutes, and supernatant and precipitates were each collected. The precipitates were subjected to the treatments of freezing-thawing and centrifugation in the same manner as described above. The supernatant was collected, and combined with the previously obtained supernatant, to give 520 ml of a crude enzyme extract.

20 Two-hundred and sixty milliliters of the crude extract was applied onto 100-ml DEAE-Sepharose FF (manufactured by Amersham Pharmacia) column equilibrated with 20 mM phosphate buffer (pH 7.0) and then non-adsorbent substances were removed by wash. Thereafter, the elution was carried out with the same buffer containing 1 M NaCl to collect 160 ml of an active fraction for 25 ceramidase. The fraction was subsequently applied onto 100-ml Phenyl-

Sephacrose FF (manufactured by Amersham Pharmacia) column equilibrated with 20 mM Tris-hydrochloric acid buffer (pH 7.5) containing 1 M NaCl. Thereafter, the elution was carried out on a concentration gradient of 2 to 0 M NaCl, and the elution was further carried out on a concentration gradient of 0 to 1%

5 Polidocanol (trade name: Lubrol PX, manufactured by Nacalai Tesque Inc.). By this chromatography, 310 ml of an active fraction for ceramidase was collected.

The resulting active fraction was applied onto 25-ml Chelating-Sephacrose FF (manufactured by Amersham Pharmacia,  $\text{Cu}^{2+}$  bound type) column equilibrated with 20 mM Tris-hydrochloric acid buffer (pH 7.5) containing  
10 0.5 M NaCl and 0.1% Lubrol PX. The column was washed with the same buffer and then with 20 mM Tris-hydrochloric acid buffer (pH 7.5) containing 0.1% Lubrol PX. Thereafter, the elution of the enzyme was carried out with 20 mM Tris-hydrochloric acid buffer (pH 7.5) containing 2 M  $\text{NH}_4\text{Cl}$  and 0.1% Lubrol PX. The eluted active fraction was concentrated by ultrafiltration, to give a  
15 concentrate. Next, the buffer in the concentrate was substituted with 20 mM Tris-hydrochloric acid buffer (pH 7.5) containing 0.1% Lubrol PX, to give an enzyme solution. Thirty milliliters of the resulting enzyme solution was further applied onto a porous HQ column ( $\phi 4.6 \times 100$  mm, manufactured by Perceptive Biosystems) equilibrated with 20 mM Tris-hydrochloric acid buffer (pH 7.5)  
20 containing 0.1% Lubrol PX. Subsequently, the elution was carried out on a concentration gradient of 0 to 0.5 M NaCl, to give an active fraction. This active fraction was applied onto hydroxyapatite column ( $\phi 7.5 \times 100$  mm, manufactured by PENTAX) equilibrated with 1 mM phosphate buffer (pH 7.0) containing 0.2 M NaCl and 0.1% Lubrol PX. The ceramidase did not adsorb to this column,  
25 and was collected on an effluent column. Thereafter, the fraction was subjected

to gel filtration chromatography by using Superose 200HR column  
( $\phi$ 10 x 300 mm, manufactured by Amersham Pharmacia) equilibrated with  
20 mM Tris-hydrochloric acid buffer (pH 7.5) containing 0.2 M NaCl and 0.3%  
Lubrol PX to give a purified ceramidase. As a result of the above purification  
5 procedures, 58 mg of a purified ceramidase preparation was obtained.

Various characteristics of the resulting purified ceramidase preparation  
were studied as described in the present specification. They were found to be as  
follows:

action: hydrolyzing ceramide to generate a sphingoid and fatty acid;

10 substrate specificity: having a substrate specificity as shown in Table 1  
listed above;

optimum pH: as shown in Figure 1, the optimum pH of this ceramidase  
being from 7.0 to 8.0;

15 temperature stability: the reduced activity not being observed when  
treated in 20 mM Tris-hydrochloric acid buffer (pH 7.5) containing 0.1%  
Polidocanol (trade name: Lubrol PX) at 37°C for 24 hours, but the activity being  
reduced to about 30% of that before the treatment when treated for 60°C for  
1 hour; and

20 molecular weight: about 94 kDa on SDS-PAGE (under reducing  
conditions); and about 73 kDa on SDS-PAGE (under reducing conditions) in this  
enzyme digested by glycopeptidase F.

#### Example 2 Partial Amino Acid Sequencing of Ceramidase

25 To 11 ml of a sample solution comprising 50 pmol ceramidase was added  
20 mM Tris-hydrochloric acid buffer (pH 7.5) containing 0.3% Lubrol PX. The

resulting sample solution was applied onto MonoQ PCl 6/5 column (100  $\mu$ l, manufactured by Amersham Pharmacia). Subsequently, the ceramidase fraction adsorbed to the column was eluted with the same buffer containing 0.4 M NaCl. By these procedures, the ceramidase-containing fraction was concentrated to a volume of 50  $\mu$ l. The resulting concentrate was subjected to SDS-polyacrylamide gel electrophoresis, and the electrophoresed gel was stained with GelCode Blue Stain reagent (manufactured by Pierce). Next, the band corresponding to the ceramidase was cut out. One-quarter of the cut-out gel fragment was subjected to extraction on 300  $\mu$ l of 0.1 M Tris-hydrochloric acid buffer (pH 9.0) containing 0.1% SDS at 37°C for 16 hours. Using the resulting extract as a sample, the N-terminal amino acid sequencing of the ceramidase was performed by using G1005A Protein Sequencing System (manufactured by Hewlett-Packard) to determine an amino acid sequence N-term. SEQ ID NO: 1 of Sequence Listing shows an amino acid sequence N-term.

In addition, the remaining three-quarter of the cut-out gel fragment was washed with 1 ml of 0.5 M Tris-hydrochloric acid buffer (pH 9.2)/50% acetonitrile at 30°C for 45 minutes. The gel was completely dried by using nitrogen gas and centrifugal concentrator, and thereafter 10  $\mu$ l of 0.5 M Tris-hydrochloric acid buffer (pH 9.2) containing 0.5  $\mu$ g of Protease Lys-C (manufactured by Wako Pure Chemical Industries, Ltd.) was added thereto. Further, 0.1 M Tris-hydrochloric acid buffer (pH 9.2) was added until the gel was completely swollen, and the mixture was kept at 37°C for 16 hours to carry out the protease digestion of the ceramidase. After the termination of the reaction, the procedures of extracting with 150  $\mu$ l of 0.1% trifluoroacetic acid / 60% acetonitrile at room temperature for 1 hour were repeated twice, and

an extract was collected. This extract was subjected to reversed phase chromatography to purify the peptide fragment. The resulting peptide fragment was analyzed by Edman degradation method using the G1005A Protein Sequencing System to determine partial amino acid sequences C-46 and C-53. SEQ ID NOs: 2 and 3 of Sequence Listing each shows an amino acid sequence for C-46 and C-53.

### Example 3 Amplification of DNA Fragment Comprising Ceramidase Gene by PCR Method

Sense mix primer 53-S1 and antisense mix primer 53-A3 were designed and synthesized with a DNA synthesizer on the basis of the partial amino acid sequence C53 of the ceramidase determined in Example 2. SEQ ID NOs: 4 and 5 of Sequence Listing each show the nucleotide sequences of the primers 53-S1 and 53-A3. PCR was carried out by using these primers. PCR was carried out with mouse liver cDNA plasmid library (manufactured by Takara Shuzo Co., Ltd.) as a template. PCR was carried out by a reaction of 94°C, 9 minutes; thereafter 40 cycles of reaction, wherein one cycle comprises a process consisting of 94°C, 0.5 minutes - 51°C, 0.5 minutes - 72°C, 1 minute; and further an incubation at 72°C for 7 minutes. By this PCR, a specific amplified DNA fragment of a size of about 70 bp was detected on agarose electrophoresis.

This amplified DNA was collected from the gel, and this DNA was incorporated in pGEM-T easy vector (manufactured by Promega) to construct a recombinant plasmid. An insert DNA fragment of the plasmid was subjected to a nucleotide sequencing. As a result, a partial nucleotide sequence P-1 of this fragment was determined. SEQ ID NO: 6 of Sequence Listing shows the

nucleotide sequence of P-1. The sequence is a sequence corresponding to the partial amino acid sequence C-53 of the ceramidase determined in Example 2. It has been confirmed that a part of the desired ceramidase gene could be obtained.

Antisense primers MA1 and MA2 were designed and synthesized on the basis of the nucleotide sequence of P-1. SEQ ID NOs: 7 and 8 of Sequence Listing each show the nucleotide sequences of the primers MA1 and MA2. In addition, sense primers T7in and T7out were designed and synthesized on the basis of the nucleotide sequence of vector pAP3neo used in the construction of the mouse liver cDNA plasmid library. SEQ ID NOs: 9 and 10 of Sequence Listing each show the nucleotide sequences of the primers T7in and T7out. Nested PCR was carried out by using these primers with the mouse liver cDNA plasmid library as a template. A 1st PCR was carried out by using the sense primer T7out and the antisense primer MA2 by a reaction at 94°C, 9 minutes; thereafter 40 cycles of reaction, wherein one cycle comprises a process consisting of 94°C, 0.5 minutes - 51°C, 0.5 minutes - 72°C, 2 minutes; and further an incubation at 72°C for 7 minutes. A 2nd PCR was carried out under the same conditions as the 1st PCR except for using the sense primer T7in and the antisense primer MA1 with a reaction mixture of the 1st PCR as a template. Consequently, an amplified DNA fragment of a size of 335 bp was obtained. This DNA fragment was used as a probe for colony hybridization described below.

#### Example 4 Cloning of Ceramidase Gene

A transformant resulting from introducing the mouse liver cDNA plasmid library was inoculated to a nylon filter (trade name Hybond-N<sup>+</sup>, manufactured by

Amersham Pharmacia) on an LB agar medium plate containing 100 µg/ml ampicillin, and about 30000 colonies were formed per one plate of 9.5 x 13.5 cm to prepare a master filter. A replica of this filter was prepared, and the resulting replica filter was respectively treated for 5 minutes on a filter paper immersed in a 10% SDS solution; 5 minutes on a filter paper immersed in a solution containing 0.5 M NaOH and 1.5 M NaCl (denaturation); 5 minutes on a filter paper immersed in 0.5 M Tris-hydrochloric acid buffer (pH 7.5) containing 3 M NaCl (neutralization); and 5 minutes on a filter paper immersed in 2 x SSC solution. Thereafter, the filter was rinsed with 2 x SSC solution. This filter was air-dried, and thereafter DNA was immobilized on a filter by ultraviolet ray irradiation to be used as a filter for colony hybridization.

As the probe for the hybridization, one prepared by <sup>32</sup>P-labeling 0.1 µg equivalent of the amplified DNA fragment obtained in Example 3 by using a DNA labeling kit, Ready To Go (manufactured by Pharmacia) in accordance with the protocol attached to the same kit. The above-mentioned filter was placed in a hybri-bag. The pre-hybridization was carried out at 60°C for 1 hour in a hybridization solution (composition: 7% PEG6000, 10% SDS solution), and thereafter the above-mentioned labeled probe was added to the mixture so as to have a concentration of 0.006 pmol/ml, and the hybridization was carried out overnight at 60°C. Next, the filter was washed three times each for 15 minutes at 60°C in the washing liquid (2 x SSC, 0.1% SDS) previously heated to 60°C. After excess water was removed from the filter, and thereafter the filter was photosensitized on an imaging plate manufactured by Fuji Photo Film for 20 minutes. Thereafter, a signal was detected with BAS1000 imaging analyzer (manufactured by Fuji Photo Film). Subsequently, colonies were collected on a

master filter corresponding to a positive signal obtained by these procedures (first screening).

The collected colonies were suspended in an LB medium containing 100 µg/ml ampicillin, and thereafter the suspension was spread over the nylon filter on an LB agar medium plate of 9.5 x 13.5 cm containing 100 µg/ml ampicillin, and about 200 to about 1000 colonies were formed per one sheet to prepare a master filter. This filter was subjected to screening of a positive clone in the same manner as the first screening, and a third screening was further carried out by the same procedures. As a result of the third screening, the positive clone which was deduced to contain the ceramidase gene could be isolated.

A plasmid was prepared from this positive clone, and this plasmid was named plasmid pLCDase. The plasmid was digested with various kinds of restriction enzymes or a combination of plural restriction enzymes. Thereafter, each of the formed DNA fragments was subcloned, and its nucleotide sequence was analyzed. By the above procedures, an entire nucleotide sequence of the DNA fragment inserted in the plasmid pLCDase was determined. The sequence is shown in SEQ ID NO: 11 of Sequence Listing. In addition, the nucleotide sequence of the open reading frame (ORF) found in the sequence and the amino acid sequence of the polypeptide encoded thereby are each shown in SEQ ID NOs: 12 and 13 of Sequence Listing. Further, the restriction enzyme map of the above-mentioned DNA fragment and the position of the open reading frame contained in the DNA fragment are shown in Figure 2.

As a result of analysis of the nucleotide sequence of the above-mentioned ORF, there were confirmed that the nucleotide sequence comprises a nucleotide

sequence encoding a partial amino acid sequence of the ceramidase elucidated by Example 2, and that this ORF encodes ceramidase. In addition, when the amino acid sequence of the ceramidase encoding the ORF was compared with the amino acid sequence of the ceramidase shown in SEQ ID NO: 1 of Sequence Listing, it was shown that the ceramidase purified from the mouse liver lacked the peptide of the N-terminal portion of the polypeptides encoded by the above-mentioned ORF. In other words, it was shown that the ceramidase was converted to a mature enzyme by going through the processing in which its N-terminal portion was removed after the translation. SEQ ID NO: 14 of Sequence Listing shows an amino acid sequence of the mature ceramidase, and SEQ ID NO: 15 of Sequence Listing shows a nucleotide sequence of the mature ceramidase, respectively.

#### Example 5 Expression of Ceramidase Gene

To CHO cells cultured in a 10% FCS-containing  $\alpha$ -MEM medium in a dish having a diameter of 35 mm ( $3 \times 10^5$  cells/dish) were added 1  $\mu$ g of the plasmid pLCDase obtained in Example 4 and 5  $\mu$ l of lipofectamine (manufactured by Life Technologies), thereby transducing a ceramidase gene into CHO cells. The cells were cultured at 37°C for 24 hours, and thereafter suspended in 100  $\mu$ l of 10 mM Tris-hydrochloric acid (pH 7.5) containing 0.1% Triton X-100, and the cells were disrupted. The determination of the activity on the ceramidase of which substrate was the above-mentioned C12-NBD-ceramide was made for the resulting solution containing disrupted cells. As a result, it was confirmed that the ceramidase in the cells intensively expressed about 1000 times more than that of the control cells in which pLCDase was not

transduced.

Further, the amount of ceramide in the cells was determined in accordance with the method described in *Analytical Biochemistry*, 244, 291-300 (1997). As a result, it was confirmed that those pLCDase-transduced cells had a significantly reduced amount of ceramide, as compared to the control cells into which pLCDase was not transduced.

#### Example 6 Cloning of Ceramidase Gene from Mouse Brain

A nitrocellulose membrane (Schleicher & Schuell, PROTRAN BA85 0.45 mm being used with a diameter of 82 mm) was placed on an LB agar medium plate containing 100 µg/ml ampicillin, and the mouse brain cDNA library (LIFE TECHNOLOGIES, SUPERScript Mouse Brain cDNA Library) was spread on each of 10 plates, so as to be about 200000 colonies per one plate, and cultured at 37°C for 10 hours. *E. coli* grown on the nitrocellulose membrane was transferred to a nylon membrane [PALL Gelman Laboratory, biodyne A diameter 82 mm (1.2 mm)], and each nylon membrane was placed on an ampicillin plate, and thereafter cultured at 37°C for 3 hours. The nitrocellulose membrane was stored at 4°C as a master filter, and the nylon membrane was placed on chloramphenicol plate, and cultured at 37°C for 16 hours. Colonies were transferred to a fresh nylon membrane from the nylon membrane, and front and back sides of the membrane were each treated for 5 minutes with 1 ml of a denaturation solution (0.5 M NaOH/1.5 M NaCl) in a state where each pair of nylon membranes was overlaid. Similarly, the nylon membranes were treated with 1 ml of a neutralization solution [0.5 M Tris-HCl (pH 7.4) / 1.5 M NaCl] for 5 minutes. The nylon membrane was peeled off, and the treated membrane was

air-dried, and thereafter baked at 80°C for 2 hours. Thereafter, the baked membrane was shaken with 200 ml of a pre-rinsing liquid [5 x SSC / 0.5% SDS / 1 ml EDTA (pH 8.0)], and disrupted *E. coli* residues were wiped off, and washed with 2 x SCC. The hybridization was carried out with 40 ml of a hybridization solution [0.5 M Church phosphate buffer/ 7% SDS / 1 mM EDTA] at 65°C for 2 hours, and thereafter the hybridization was carried out at 65°C for 16 hours in 40 ml of the hybridization solution containing the denatured probe. As the probe, an EcoRI-EcoRI fragment of a size of 2.7 kbp of the plasmid pAPLCD carrying mouse ceramidase gene was used. After the termination of the hybridization, washing with 100 ml of a washing liquid (40 mM Church phosphate buffer/ 1% SDS) at 65°C for 15 minutes was carried out twice. Further, washing with 100 ml of a high stringent washing liquid (0.2 x SSC / 0.1% SDS) at 65°C for 15 minutes was carried out. The membrane was air-dried, and thereafter exposed on an IP-plate for 1 hour, and analyzed with BAS 1500. The positive part of the nitrocellulose membrane was cut out in a diameter of about 6 mm, and suspended in 1 ml of the LB medium. Thereafter, a 200 ml sample prepared by diluting the positive part 4000-folds was spread over an ampicillin-containing LB plate, and a 2nd screening was carried out.

Similarly, a 200 ml sample prepared by diluting the positive part 10000-folds was spread over an ampicillin-containing LB plate, to prepare a library. A 3rd screening was carried out by using this library. The isolated clone (pSB CD) was subcloned, and thereafter the nucleotide sequence was determined by a conventional method. This sequence is shown in SEQ ID NO: 16.

In the above-mentioned sequence, an ORF having the identical sequence as the ceramidase gene derived from mouse liver described in Example 4 was

found. Incidentally, the sequences of 5' non-translational region and 3' non-translational region were different from those derived from the mouse liver. In addition, the isolated plasmid pSBCD was transduced into CHO cells in the same manner as in Example 5. As a result, it was found that the ceramidase was expressed in the cells.

#### Example 7 Genomic Cloning of Human Ceramidase Gene

Sense primer U1107 having the sequence of SEQ ID NO: 17 of Sequence Listing and antisense primer L1311 having the sequence of SEQ ID NO: 18 of Sequence Listing were synthesized on the basis of the sequence of the ceramidase gene derived from mouse liver determined in Example 4. The sequence of U1107 primer corresponds to a sequence consisting of base nos.: 1107 - 1130 of SEQ ID NO: 12 of Sequence Listing, and the sequence of L1311 primer corresponds to a nucleotide sequence complementary to a sequence consisting of base nos.: 1311 - 1334 of SEQ ID NO: 12 of Sequence Listing.

Genomic DNA was purified from human hepatoma cell Huh7 by a conventional method. PCR was carried out by using the U1107 primer and the L1311 primer with 625 ng of the resulting genomic DNA as a template. PCR was carried out by a reaction at 94°C, 9 minutes; thereafter 40 cycles of reaction, wherein one cycle comprises a process consisting of 94°C, 0.5 minutes - 55°C, 0.5 minutes - 72°C, 3 minutes; and further an incubation at 72°C for 7 minutes. Thereafter, the resulting reaction product was subjected to agarose electrophoresis. As a result, it was confirmed that a DNA fragment of a size of about 2 kbp was amplified by the PCR.

This DNA fragment was collected from the gel by using Sephaglas (manufactured by Pharmacia), and the resulting fragment was incorporated in pGEM-T easy vector (manufactured by Promega) to construct a recombinant plasmid. Next, the nucleotide sequence of the DNA fragment insert of the resulting plasmid was determined. The above-mentioned sequence was a sequence corresponding to 96289 - 98478 of the Accession No. AC012131 Complement registered in the GenBank Data Base. In addition, the amino acid sequence encoded by the above-mentioned sequence was analyzed. As a result, a region encoding an amino acid sequence showing homology to a region of the amino acid nos.: 370 - 444 of the amino acid sequence of the ceramidase derived from mouse liver of SEQ ID NO: 4 of Sequence Listing and a region encoding an amino acid sequence showing homology were found.

Incidentally, AC012131 also has homology with the ceramidase gene derived from mouse liver determined in Example 4.

#### SEQUENCE LISTING FREE TEXT

In the amino acid sequence of SEQ ID NO: 1, each of Xaa in the amino acid numbers: 7, 9 and 13 stands for an unknown amino acid.

SEQ ID NO: 4 is a sequence of synthetic oligonucleotide primers. In the above-mentioned sequence, each of n in the base numbers: 6, 9 and 15 stands for G, A, T or C.

SEQ ID NO: 5 is a sequence of synthetic oligonucleotide primers. In the above-mentioned sequence, each of n in the base numbers: 3, 6 and 15 stands for G, A, T or C.

SEQ ID NO: 7 is a sequence of synthetic oligonucleotide primers.

SEQ ID NO: 8 is a sequence of synthetic oligonucleotide primers.

SEQ ID NO: 9 is a sequence of synthetic oligonucleotide primers.

SEQ ID NO: 10 is a sequence of synthetic oligonucleotide primers.

SEQ ID NO: 11 is a sequence of synthetic oligonucleotide primers.

5 SEQ ID NO: 17 is a sequence of synthetic oligonucleotide primers.

SEQ ID NO: 18 is a sequence of synthetic oligonucleotide primers.

#### INDUSTRIAL APPLICABILITY

10 According to the present invention, a gene encoding a neutral/alkaline  
ceramidase derived from a mammal is provided, and a genetic engineering  
method for producing a ceramidase using the gene. Also, the oligonucleotide  
probe and primer of the present invention are useful for detection of the above-  
mentioned gene, and can be applied to studies on *in vivo* ceramide metabolism.  
15 Further, according to the present invention, an antisense nucleic acid (DNA,  
RNA) of the gene of the present invention is provided. The gene and its  
antisense nucleic acid are useful in the control of the ceramidase activity and in  
the regulations of the *in vivo* ceramide metabolism system. Therefore, a method  
of regulating an amount of ceramide which can be applied to treatments of  
diseases caused by the abnormal amount of ceramide is provided.

20

## CLAIMS

1. A gene having a nucleotide sequence of a nucleic acid selected from the group consisting of:

- 5 (A) a nucleic acid encoding a polypeptide having the amino acid sequence of SEQ ID NO: 14 of the Sequence Listing or a partial sequence thereof, the polypeptide possessing a ceramidase activity;
- 10 (B) a nucleic acid having a nucleotide sequence of SEQ ID NO: 15 of the Sequence Listing or a partial sequence thereof and encoding a polypeptide possessing a ceramidase activity;
- (C) a nucleic acid encoding a polypeptide consisting of an amino acid sequence resulting from deletion, addition, insertion or substitution of at least one amino acid residue in the amino acid sequence of SEQ ID NO: 14 of the Sequence Listing, the polypeptide possessing a ceramidase activity;
- 15 (D) a nucleic acid consisting of a nucleotide sequence resulting from deletion, addition, insertion or substitution of at least one base in the nucleotide sequence of SEQ ID NO: 15 of the Sequence Listing and encoding a polypeptide possessing a ceramidase activity;
- 20 (E) a nucleic acid capable of hybridizing to a complementary strand of a nucleic acid of any one of the above (A) to (D), under stringent conditions, and encoding a polypeptide possessing a ceramidase activity; and
- (F) a nucleic acid having a nucleotide sequence different from the nucleic acid of any one of above (A) to (E) via degeneracy and encoding a polypeptide possessing a ceramidase activity.
- 25

0937521.092601

2. The gene according to claim 1, wherein the ceramidase activity of the polypeptide is capable of being detected by the following steps:

- (a) incubating a gene expression product in a reaction mixture [composition: 550 pmol of C12-NBD-ceramide and 1.0% (W/V) sodium cholate in 20  $\mu$ l of 25 mM Tris-hydrochloric acid buffer (pH 7.5)] at 37°C for 30 minutes, to react the mixture; and
- (b) detecting generation of a C12-NBD-fatty acid in the reaction product.

3. The gene according to claim 1 or 2, wherein the polypeptide exhibits at least the following characteristics:

- (i) action: hydrolyzing ceramide, to generate sphingoid and a fatty acid;
- (ii) substrate specificity: hydrolyzing N-acylsphingosine; but not acting on galactosylceramide, sulfatide, GM1a, and sphingomyelin;
- (iii) optimum pH: being from 7.0 to 8.0; and
- (iv) lowering of the activity not being found when treated in 20 mM Tris-hydrochloric acid (pH 7.5) containing 0.1% polidocanol at 37°C for 24 hours, but the activity lowering by a treatment at 60°C for 1 hour to about 30% of the activity before the treatment.

4. A recombinant DNA comprising the gene of any one of claims 1 to 3.

5. An expression vector for a microorganism, an animal cell or a plant cell, comprising the gene of any one of claims 1 to 3 or the recombinant DNA of claim 4.

6. A transformant carrying the expression vector of claim 5.
7. A method for producing a polypeptide possessing a ceramidase activity, characterized by culturing the transformant of claim 6 under conditions appropriate for expression of the ceramidase gene and production of the polypeptide encoded by the gene, and collecting a polypeptide possessing a ceramidase activity from the resulting culture.
8. A polypeptide having the amino acid sequence of SEQ ID NO: 14 of the Sequence Listing or a partial sequence thereof and possessing a ceramidase activity.
9. A polypeptide possessing a ceramidase activity, encoded by the gene of any one of claims 1 to 3.
10. The polypeptide according to claim 8 or 9, wherein the ceramidase activity is capable of being detected by the following steps:
- (a) incubating a gene expression product in a reaction mixture [composition: 550 pmol of C12-NBD-ceramide and 1.0% (W/V) sodium cholate in 20  $\mu$ l of 25 mM Tris-hydrochloric acid buffer (pH 7.5)] at 37°C for 30 minutes, to react the mixture; and
  - (b) detecting generation of a C12-NBD-fatty acid in the reaction product.
11. An antisense DNA which is complementary to the gene of any one of claims 1 to 3 or a part thereof.

12. An antisense RNA which is complementary to the gene of any one of claims 1 to 3 or a part thereof.

5 13. An expression vector comprising the antisense DNA of claim 11.

14. An oligonucleotide probe or primer, capable of specifically hybridizing to the gene of any one of claims 1 to 3 or a complementary strand thereof.

10 15. An antibody or a fragment thereof, capable of specifically binding to the polypeptide of any one of claims 8 to 10.

15 16. A method for detecting a gene encoding a polypeptide possessing a ceramidase activity, comprising using the oligonucleotide probe or primer of claim 14.

17. A kit for the use in detection of a gene encoding a polypeptide possessing a ceramidase activity, comprising the oligonucleotide probe and/or primer of claim 14.


20 18. A method for detecting a polypeptide possessing a ceramidase activity, comprising using the antibody or a fragment thereof of claim 15.

25 19. A kit for the use in detection of a polypeptide possessing a ceramidase activity, comprising the antibody or a fragment thereof of claim 15.

20. A method of controlling an amount of ceramide in a cell and/or in a tissue, characterized by introducing the gene of any one of claims 1 to 3 or an antisense nucleic acid thereof into the cell and/or the tissue, thereby controlling the amount of ceramide in the cell and/or in the tissue.
- 5


0997521 09E01  
T09E01 T254E660

FIG. 1



T09260\* T2526650

FIG. 2



PLEASE NOTE:  
YOU MUST  
COMPLETE THE  
FOLLOWING: ↓COMBINED DECLARATION AND POWER OF ATTORNEY  
FOR PATENT AND DESIGN APPLICATIONS

As a below named inventor, I hereby declare that: my residence, post office address and citizenship are as stated next to my name; that I verily believe that I am the original, first and sole inventor (if only one inventor is named below) or an original, first and joint inventor (if plural inventors are named below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

CERAMIDASE GENE

Insert Title: →

Fill in Appropriate  
Information —

For Use →

Without

Specification

Attached:

the specification of which is attached hereto. If not attached hereto,

the specification was filed on \_\_\_\_\_ as

United States Application Number \_\_\_\_\_;

and amended on \_\_\_\_\_ (if applicable); and/or

the specification was filed on March 24, 2000 as PCTInternational Application Number PCT/JP00/01802; and was

amended under PCT Article 19 on \_\_\_\_\_ (if applicable)

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, §1.56.

I do not know and do not believe the same was ever known or used in the United States of America before my or our invention thereof, or patented or described in any printed publication in any country before my or our invention thereof or more than one year prior to this application, that the same was not in public use or on sale in the United States of America more than one year prior to this application, that the invention has not been patented or made the subject of an inventor's certificate issued before the date of this application in any country foreign to the United States of America on an application filed by me or my legal representatives or assigns more than twelve months (six months for designs) prior to this application, and that no application for patent or inventor's certificate on this invention has been filed in any country foreign to the United States of America prior to this application by me or my legal representatives or assigns, except as follows.

I hereby claim foreign priority benefits under Title 35, United States Code, §119 (a)-(d) of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed:

## Prior Foreign Application(s)

## Priority Claimed

Insert Priority  
Information: →  
(if appropriate)

|                             |                           |   |   |                             |
|-----------------------------|---------------------------|---|---|-----------------------------|
| <u>11-84743</u><br>(Number) | <u>Japan</u><br>(Country) | <u>March 26, 1999</u><br>(Month / Day / Year Filed) | <input checked="" type="checkbox"/> Yes | <input type="checkbox"/> No |
| _____<br>(Number)           | _____<br>(Country)        | _____<br>(Month / Day / Year Filed)                 | <input type="checkbox"/> Yes            | <input type="checkbox"/> No |
| _____<br>(Number)           | _____<br>(Country)        | _____<br>(Month / Day / Year Filed)                 | <input type="checkbox"/> Yes            | <input type="checkbox"/> No |
| _____<br>(Number)           | _____<br>(Country)        | _____<br>(Month / Day / Year Filed)                 | <input type="checkbox"/> Yes            | <input type="checkbox"/> No |

I hereby claim the benefit under Title 35, United States Code, §119(e) of any United States provisional application(s) listed below.

Insert Provisional  
Application(s): →  
(if any)

|                               |                        |
|-------------------------------|------------------------|
| _____<br>(Application Number) | _____<br>(Filing Date) |
| _____<br>(Application Number) | _____<br>(Filing Date) |

All Foreign Applications, if any, for any Patent or Inventor's Certificate Filed More than 12 Months (6 Months for Designs) Prior to the Filing Date of This Application:

Insert Requested  
Information: →  
(if appropriate)

| Country | Application Number | Date of Filing (Month / Day / Year) |
|---------|--------------------|-------------------------------------|
| _____   | _____              | _____                               |
| _____   | _____              | _____                               |

I hereby claim the benefit under Title 35, United States Code, §120 of any United States and/or PCT application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States and/or PCT application in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, §1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application:

Insert Prior U.S.  
Application(s): →  
(if any)

|                               |                        |  |
|-------------------------------|------------------------|--|
| _____<br>(Application Number) | _____<br>(Filing Date) | _____<br>(Status — patented, pending, abandoned) |
| _____<br>(Application Number) | _____<br>(Filing Date) | _____<br>(Status — patented, pending, abandoned) |

I hereby appoint the following attorneys to prosecute this application and/or an international application based on this application and to transact all business in the Patent and Trademark Office connected therewith and in connection with the resulting patent based on instructions received from the entity who first sent the application papers to the attorneys identified below, unless the inventor(s) or assignee provides said attorneys with a written notice to the contrary:

1422-0493P

|                     |                   |                       |                   |
|---------------------|-------------------|-----------------------|-------------------|
| Raymond C. Stewart  | (Reg. No. 21,066) | Terrell C. Birch      | (Reg. No. 19,382) |
| Joseph A. Kolasch   | (Reg. No. 22,463) | James M. Slattery     | (Reg. No. 28,380) |
| Bernard L. Sweeney  | (Reg. No. 24,448) | Michael K. Mutter     | (Reg. No. 29,680) |
| Charles Gorenstein  | (Reg. No. 29,271) | Gerald M. Murphy, Jr. | (Reg. No. 28,977) |
| Leonard R. Svensson | (Reg. No. 30,330) | Terry L. Clark        | (Reg. No. 32,644) |
| Andrew D. Meikle    | (Reg. No. 32,868) | Marc S. Weiner        | (Reg. No. 32,181) |
| Joe McKinney Muncy  | (Reg. No. 32,334) | C. Joseph Faraci      | (Reg. No. 32,350) |
| Donald J. Daley     | (Reg. No. 34,313) | John W. Bailey        | (Reg. No. 32,881) |
| John A. Castellano  | (Reg. No. 35,094) |                       |                   |

Send Correspondence to: **BIRCH, STEWART, KOLASCH & BIRCH, LLP**  
P.O. Box 747 • Falls Church, Virginia 22040-0747  
Telephone: (703) 205-8000 • Facsimile: (703) 205-8050

PLEASE NOTE:  
YOU MUST  
COMPLETE  
THE  
FOLLOWING:

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full Name of First or  
Sole Inventor  
Insert Name of  
Inventor  
Insert Date This  
Document is Signed

Insert Residence  
Insert Citizenship

Insert Post Office  
Address

Full Name of Second  
Inventor, if any,  
see above

Full Name of Third  
Inventor, if any,  
see above

Full Name of Fourth  
Inventor, if any,  
see above

Full Name of Fifth  
Inventor, if any,  
see above

|   |  |             |  |                      |  |             |  |
|---|--|-------------|--|----------------------|--|-------------|--|
| GIVEN NAME  |  | FAMILY NAME |  | INVENTOR'S SIGNATURE |  | DATE*       |  |
| Makoto  |  | ITO         |  | Makoto Ito           |  | Aug. 28, 01 |  |
| Residence (City, State & Country)   |  |             |  |                      |  | CITIZENSHIP |  |
| Fukuoka-shi, Fukuoka 819-0013 Japan Japan JPX                                 |  |             |  |                      |  | Japan       |  |
| POST OFFICE ADDRESS (Complete Street Address including City, State & Country) |  |             |  |                      |  |             |  |
| 34-1, Atagohama 4-chome, Nishi-ku, Fukuoka-shi, Fukuoka 819-0013 Japan        |  |             |  |                      |  |             |  |
| GIVEN NAME  |  | FAMILY NAME |  | INVENTOR'S SIGNATURE |  | DATE*       |  |
|   |  |             |  |                      |  |             |  |
| Residence (City, State & Country)   |  |             |  |                      |  | CITIZENSHIP |  |
|   |  |             |  |                      |  |             |  |
| POST OFFICE ADDRESS (Complete Street Address including City, State & Country) |  |             |  |                      |  |             |  |
|   |  |             |  |                      |  |             |  |
| GIVEN NAME  |  | FAMILY NAME |  | INVENTOR'S SIGNATURE |  | DATE*       |  |
|   |  |             |  |                      |  |             |  |
| Residence (City, State & Country)   |  |             |  |                      |  | CITIZENSHIP |  |
|   |  |             |  |                      |  |             |  |
| POST OFFICE ADDRESS (Complete Street Address including City, State & Country) |  |             |  |                      |  |             |  |
|   |  |             |  |                      |  |             |  |
| GIVEN NAME  |  | FAMILY NAME |  | INVENTOR'S SIGNATURE |  | DATE*       |  |
|   |  |             |  |                      |  |             |  |
| Residence (City, State & Country)   |  |             |  |                      |  | CITIZENSHIP |  |
|   |  |             |  |                      |  |             |  |
| POST OFFICE ADDRESS (Complete Street Address including City, State & Country) |  |             |  |                      |  |             |  |
|   |  |             |  |                      |  |             |  |
| GIVEN NAME  |  | FAMILY NAME |  | INVENTOR'S SIGNATURE |  | DATE*       |  |
|   |  |             |  |                      |  |             |  |
| Residence (City, State & Country)   |  |             |  |                      |  | CITIZENSHIP |  |
|   |  |             |  |                      |  |             |  |
| POST OFFICE ADDRESS (Complete Street Address including City, State & Country) |  |             |  |                      |  |             |  |
|   |  |             |  |                      |  |             |  |

## SEQUENCE LISTING

<110> Takara Shuzo Co., Ltd.

<120> A gene encoding ceramidase

<130> 00-011-PCT

<140> JP 11/84743

<141> 1999-3-26

<160> 18

<210> 1

<211> 21

<212> PRT

<213> Mouse

<220>

<222> 7, 9, 13

<223> Xaa is an unknown amino acid.

<400> 1

Phe Ser Gly Tyr Tyr Ile Xaa Val Xaa Arg Ala Asp Xaa Thr Gly

1

5

10

15

Lys Val Asn Asp Ile Asn

20

<210> 2

T09250" T23/0660

<211> 10

<212> PRT

<213> Mouse

<220>

<222> 9

<223> Xaa is an unknown amino acid.

<400> 2

Ala Ile Ala Thr Asp Thr Val Ala Xaa Met

1 5 10

<210> 3

<211> 35

<212> PRT

<213> Mouse

<220>

<222> 29, 30

<223> Xaa is an unknown amino acid.

<400> 3

Gly Tyr Leu Pro Gly Gln Gly Pro Phe Val Asn Gly Phe Ala Ser

1 5 10 15

Ser Asn Leu Gly Asp Val Ser Pro Asn Ile Leu Gly Pro Xaa Xaa

20 25 30

Val Asn Thr Gly Glu

35

09/937521

<210> 4

<211> 17

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthesized oligonucleotide for primer.

<220>

<222> 6, 9, 15

<223> "n" is G or A or T or C.

<400> 4

carggncnt tygtngc

17

<210> 5

<211> 17

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthesized oligonucleotide for primer.

<220>

<222> 3, 6, 15

<223> "n" is G or A or T or C.

<400> 5

ggncnagda trttngg

17

09/937521

<210> 6

<211> 38

<212> DNA

<213> Mouse

<400> 6

gcaggctttg cttcatcaaa tctcgagac gtgtcacc

38

<210> 7

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthesized oligonucleotide for primer.

<400> 7

ttgatgaagc aaagcctgc

19

<210> 8

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthesized oligonucleotide for primer

<400> 8

09937521.05601

ggtgacacgt ctccgagat

19

<210> 9

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthesized oligonucleotide for primer

<400> 9

taatacgact cactataggg

20

<210> 10

<211> 17

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthesized oligonucleotide for primer

<400> 10

tctgctctaa aagctgc

17

<210> 11

<211> 3108

<212> DNA

<213> Mouse

0937521-09260

&lt;400&gt; 11

cctgcgccac ttctctctcc cggtcaatc gggagcctt ttctctcccc cgtctgccg 60  
 ctgccgccat ctccaccctt gcctgcccc ggggtctgtg gacgcccgagg cagagagcaa 120  
 gcaccgagct gggcctgtct gagaccggag accagcggcc cgcccccccg cccgctgcga 180  
 gcctctctgag cagctccgga acagcttact ttctgtttcc atctctttcg gaccgggttg 240  
 gcctctccaa aagccacttc tcctaactct tatcaagggtt caaaggctaa aggtctgtac 300  
 acatgagtgc tgggtgtgctt agaggcatcg ggtccctttc agctggagtt gcagtacttg 360  
 tgagtgccat ggaatccaaa ttcggcaaga gatacaatct aaactctcaa ctactccaga 420  
 ttcaagggtc acctcacttt ctggttacca aaggagcttt gcggggccgc tctgacatcc 480  
 agtagatttg gaaacacatt gagaaatcag cctgagcaac ctgcaaggca caaggcacia 540  
 gattctgcat ggttatttgc tctcccagga ggtgaacact tgttttgatt cacagagtca 600  
 ggggttgagat gccagttgt tctcatctt ggctcagaag aagcacctag gaataaaagc 660  
 tctaagctgg tattaagtag aatgggctta aagtccacta caggaaacaa cagctagtga 720  
 cagaaatggc aaagcgaacc ttctccacct tggaggcatt cctcattttc cttctggtaa 780  
 taatgacagt catcacagt gcccttctca cctcttgtt tgttaccagt gggaccattg 840  
 aaaaccacaa agattcagga aatcacttgt tttcaaccac tctgggctcc acgacaacc 900  
 agccccctcc aattacacag actccaaact tcccttcatt tcggaacttc agtggctact 960  
 acattggcgt tgggagagcg gattgcacag gacaagtgtc agatatcaat ttgatgggct 1020  
 atggcaaaaa tggccagaat gcacggggtc tctcaccag gctgttcagc cgtgctttta 1080  
 tcttggcgga tccagatggg tcaaatcgaa tggcatttgt gagcgtggaa ctatgtatga 1140  
 tttccaacg actgaggttg gaggtcctga agagactaga gagtaaataat ggctctctgt 1200  
 atcgaagaga caatgttatc ctgagtcca ttacacaca ctctggccca gcagggtttt 1260  
 tccaatatac actctatata ctgccagcg agggattcag caaccggacc tttcagtaca 1320  
 tagtctctgg gatcatgaag agcattgata tagctcacac aaatcttaaa ccaggcaaaa 1380  
 tctttatcaa caaaggaaat gttgctaag tgcagatcaa ccgaagcccc tctcttacc 1440  
 ttctgaatcc acagtcatg agagcaaggt attcttcaa cacagacaag gaaatgctgg 1500  
 tcttgaaact ggtggatttg aatggagaag acttgggtct tatcagctgg ttgccaatcc 1620  
 acccgtgag catgaacaat agcaaccact ttgtgaatag tgacaatatg ggctatgcgg 1680  
 cttacctttt tgagcaagaa aagaacaaag gctatctgcc tggacaggga ccgtttgtag 1680

0937521.092501

caggctttgc ttcacaaat ctccgagacg tgcacccaa cattcttggc ccgcattgtg 1740  
 tcaacacagg ggagtcttgt gacaacgaca agagcacctg tccaacggt gggcctagca 1800  
 tgtgcatggc cagcggacct ggacaagaca tgtttgagag cacacacatt ataggacgga 1860  
 tcattctatca gaaggccaag gagctgtatg cctctgcctc ccaggagggtg accggcccag 1920  
 tgcttgacg tcaccagtgg gtgaacatga cagatgtgag cgtccagctc aatgccacac 1980  
 acacagtga gacgtgtaaa cctgccctgg gctacagttt tgccgcaggc acaattgatg 2040  
 gagtttcggg cctcaatatt acacagggaa ctacggaagg ggatccattc tgggacactc 2100  
 ttcgggacca gctcttggga aaaccatctg aagagattgt agagtgtcag aaacccaaac 2160  
 caatcctgct tcacagtga gagctgacga taccacatcc ttggcaacca gatattgttg 2220  
 atgttcagat tgttaccgtt gggctccttg ccatagctgc tatccctggg gaattaacaa 2280  
 ccatgtcggg acgaagattt cgtgaggcaa ttaaaaaaga atttgactt tatgggatga 2340  
 aggatatgac cgttggtatc gcagggtctaa gcaatgttta tacacattac attaccacat 2400  
 atgaagaata ccaggctcag cggtagcagg cagcatctac aatctatgga ccacacaccc 2460  
 tgtctgcata catccaactc ttcagagacc ttgctaaggc aattgctacg gacacagtag 2520  
 ccaacatgag cagtggctcc gagcctccat tctcaaaaa tctaatactg tcaattattc 2580  
 ctaatattgc ggatagagca ccaattggca aacatttttg ggatgtcttg cagccagcaa 2640  
 aacctgaata cagagtggga gaagtgggtg aagttatatt ttaggcgct aacccaaaga 2700  
 attcagcaga gaaccagacc catcaaacct tctcactgt ggagaaatac gaggactctg 2760  
 tagctgactg gcagataatg tataacgatg cctcctggga gacgaggttt tattggcaca 2820  
 aaggaatact gggctcagc aatgcaacaa tatactggca tattccagat actgcctacc 2880  
 ctggaatcta cagaataaga tattttggac acaatcgga gcaggaactt ctgaaacccg 2940  
 ctgtcatact agcatttgaa ggaatttctt ctccttttga agttgtcact acttagtgaa 3000  
 aagttgacag atattgaaga aaagcttttc tctgtgcaca ttatagagtg aattcacaaa 3060  
 aaaaaaaaa aaaaaaaaa aaaaaaaaa aaaaaaaaa aaaaaaaaa 3108

<210> 12

<211> 2271

<212> DNA

<213> Mouse

0937521.09501

&lt;400&gt; 12

atggcaaagc gaaccttctc caccttggag gcattcctca ttttcttct ggtaataatg 60  
 acagtcatca cagtggccct tctcacctc ttgtttgta ccagtgggac cattgaaaac 120  
 cacaagatt caggaaatca ctgttttca accactctgg gctccacgac aaccagccc 180  
 cctccaatta cacagactcc aaacttcctt tcatttcgga acttcagtgg ctactacatt 240  
 ggcgttggga gagcggattg cacaggacaa gtgtcagata tcaatttgat ggcctatggc 300  
 aaaaatggcc agaatgcacg gggctcctc accaggctgt tcagccgtgc ttttatcttg 360  
 gcgatccag atgggtcaaa tcgaatggca tttgtgagcg tggaactatg tatgatttcc 420  
 caacgactga ggttggaggt cctgaagaga ctagagagta aatatggctc tctgtatcga 480  
 agagacaatg ttatcctgag tgccattcac acacactctg gccagcagg gtttttcaa 540  
 tatacactct atatactgc cagcaggga ttcagcaacc ggaccttca gtacatagtc 600  
 tctgggatca tgaagagcat tgatatagct cacacaaatc ttaaaccagg caaaatcttt 660  
 atcaacaaag gaaatgttgc taatgtgcag atcaaccgaa gcccctcctc ttaccttctg 720  
 aatccacagt cagagagagc aaggtattct tcaaacacag acaaggaaat gctggcttgg 780  
 aaactggigg atttgaatgg agaagacttg ggtcttatca gctggtttgc catccacccc 840  
 gtgagcatga acaatagcaa ccactttgtg aatagtgaac atatgggcta tgcggcttac 900  
 ctttttgagc aagaaaagaa caaaggctat ctgcctggac agggaccgtt tgtagcaggc 960  
 ttgtcttcat caaatctcgg agacgtgtca cccaacattc ttggcccgca ttgtgtcaac 1020  
 acaggggagt cttgtgacaa cgacaagagc acctgtccca acggtgggccc tagcatgtgc 1080  
 atggccagcg gacctggaca agacatgttt gagagcacac acattatagg acggatcacc 1140  
 tatcagaagg ccaaggagct gtatgcctct gcctcccagg aggtgaccgg cccagtgttt 1200  
 gcagctcacc agtgggtgaa catgacagat gtgagcgtcc agctcaatgc cacacacaca 1260  
 gtgaagacgt gtaaacctgc cctgggctac agttttgccg caggcacaat tgatggagtt 1320  
 tcgggcctca atattacaca gggaactacg gaaggggac cttctggga cactcttcgg 1380  
 gaccagctct tgggaaaacc atctgaagag attgtagagt gtcagaaacc caaaccaatc 1440  
 ctgcttcaca gtggagagct gacgatacca catccttggc aaccagatat tgttgatgtt 1500  
 cagattgita ccgttgggtc cttggccata gctgctatcc ctggggaatt aacaaccatg 1620  
 tcgggacgaa gatttcgtga ggcaattaaa aaagaatttg cactttatgg gatgaaggat 1620

0937521.09360

atgaccgttg ttatcgagg tctaagcaat gtttatacac attacattac cacatatgaa 1680  
 gaataccagg ctacgaggta cgaggcagca tctacaatct atggaccaca caccctgtct 1740  
 gcatacatcc aactcttcag agaccttgct aaggcaattg ctacggacac agtagccaac 1800  
 atgagcagtg gtcccagacc tccattcttc aaaaatctaa tagcttcact ttttctaata 1860  
 attgaggata gagcaccaat tggcaaacaat tttggggatg tcttgagacc agcaaaacct 1920  
 gaatacagag tgggagaagt ggttgaagtt atattttagt gcgctaaccc aaagaattca 1980  
 gcagagaacc agacccatca aaccttctc actgtggaga aatacgagga ctctgtagct 2040  
 gactggcaga taatgtataa cgatgcctcc tgggagacga ggttttattg gcacaaagga 2100  
 atactgggtc tgagcaatgc aacaatatac tggcatattc cagatactgc ctaccctgga 2160  
 atctacagaa taagatattt tggacacaat cggaagcagg aacttctgaa acccgctgtc 2220  
 atactagcat ttgaaggaat ttcttctcct tttgaagttg tcaactacta g 2271

<210> 13

<211> 756

<212> PRT

<213> Mouse

<400> 13

Met Ala Lys Arg Thr Phe Ser Thr Leu Glu Ala Phe Leu Ile Phe

1 5 10 15

Leu Leu Val Ile Met Thr Val Ile Thr Val Ala Leu Leu Thr Leu

20 25 30

Leu Phe Val Thr Ser Gly Thr Ile Glu Asn His Lys Asp Ser Gly

35 40 45

Asn His Trp Phe Ser Thr Thr Leu Gly Ser Thr Thr Thr Gln Pro

50 55 60

Pro Pro Ile Thr Gln Thr Pro Asn Phe Pro Ser Phe Arg Asn Phe

65 70 75

Ser Gly Tyr Tyr Ile Gly Val Gly Arg Ala Asp Cys Thr Gly Gln

0937521.0937521.0937521

|   |     |     |
|---|-----|-----|
| 80  | 85  | 90  |
| Val Ser Asp Ile Asn Leu Met Gly Tyr Gly Lys Asn Gly Gln Asn |     |     |
| 95  | 100 | 105 |
| Ala Arg Gly Leu Leu Thr Arg Leu Phe Ser Arg Ala Phe Ile Leu |     |     |
| 110   | 115 | 120 |
| Ala Asp Pro Asp Gly Ser Asn Arg Met Ala Phe Val Ser Val Glu |     |     |
| 125   | 130 | 135 |
| Leu Cys Met Ile Ser Gln Arg Leu Arg Leu Glu Val Leu Lys Arg |     |     |
| 140   | 145 | 150 |
| Leu Glu Ser Lys Tyr Gly Ser Leu Tyr Arg Arg Asp Asn Val Ile |     |     |
| 155   | 160 | 165 |
| Leu Ser Ala Ile His Thr His Ser Gly Pro Ala Gly Phe Phe Gln |     |     |
| 170   | 175 | 180 |
| Tyr Thr Leu Tyr Ile Leu Ala Ser Glu Gly Phe Ser Asn Arg Thr |     |     |
| 185   | 190 | 195 |
| Phe Gln Tyr Ile Val Ser Gly Ile Met Lys Ser Ile Asp Ile Ala |     |     |
| 200   | 205 | 210 |
| His Thr Asn Leu Lys Pro Gly Lys Ile Phe Ile Asn Lys Gly Asn |     |     |
| 215   | 220 | 225 |
| Val Ala Asn Val Gln Ile Asn Arg Ser Pro Ser Ser Tyr Leu Leu |     |     |
| 230   | 235 | 240 |
| Asn Pro Gln Ser Glu Arg Ala Arg Tyr Ser Ser Asn Thr Asp Lys |     |     |
| 245   | 250 | 255 |
| Glu Met Leu Val Leu Lys Leu Val Asp Leu Asn Gly Glu Asp Leu |     |     |
| 260   | 265 | 270 |
| Gly Leu Ile Ser Trp Phe Ala Ile His Pro Val Ser Met Asn Asn |     |     |
| 275   | 280 | 285 |
| Ser Asn His Phe Val Asn Ser Asp Asn Met Gly Tyr Ala Ala Tyr |     |     |
| 290   | 295 | 300 |

0937521.09601

|   |     |     |     |
|---|-----|-----|-----|
| Leu Phe Glu Gln Glu Lys Asn Lys Gly Tyr Leu Pro Gly Gln Gly | 305 | 310 | 315 |
| Pro Phe Val Ala Gly Phe Ala Ser Ser Asn Leu Gly Asp Val Ser | 320 | 325 | 330 |
| Pro Asn Ile Leu Gly Pro His Cys Val Asn Thr Gly Glu Ser Cys | 335 | 340 | 345 |
| Asp Asn Asp Lys Ser Thr Cys Pro Asn Gly Gly Pro Ser Met Cys | 350 | 355 | 360 |
| Met Ala Ser Gly Pro Gly Gln Asp Met Phe Glu Ser Thr His Ile | 365 | 370 | 375 |
| Ile Gly Arg Ile Ile Tyr Gln Lys Ala Lys Glu Leu Tyr Ala Ser | 380 | 385 | 390 |
| Ala Ser Gln Glu Val Thr Gly Pro Val Leu Ala Ala His Gln Trp | 395 | 400 | 405 |
| Val Asn Met Thr Asp Val Ser Val Gln Leu Asn Ala Thr His Thr | 410 | 415 | 420 |
| Val Lys Thr Cys Lys Pro Ala Leu Gly Tyr Ser Phe Ala Ala Gly | 425 | 430 | 435 |
| Thr Ile Asp Gly Val Ser Gly Leu Asn Ile Thr Gln Gly Thr Thr | 440 | 445 | 450 |
| Glu Gly Asp Pro Phe Trp Asp Thr Leu Arg Asp Gln Leu Leu Gly | 455 | 460 | 465 |
| Lys Pro Ser Glu Glu Ile Val Glu Cys Gln Lys Pro Lys Pro Ile | 470 | 475 | 480 |
| Leu Leu His Ser Gly Glu Leu Thr Ile Pro His Pro Trp Gln Pro | 485 | 490 | 495 |
| Asp Ile Val Asp Val Gln Ile Val Thr Val Gly Ser Leu Ala Ile | 500 | 505 | 510 |
| Ala Ala Ile Pro Gly Glu Leu Thr Thr Met Ser Gly Arg Arg Phe |     |     |     |

09/937521-092601

|   |     |     |
|---|-----|-----|
| 515   | 520 | 525 |
| Arg Glu Ala Ile Lys Lys Glu Phe Ala Leu Tyr Gly Met Lys Asp |     |     |
| 530   | 535 | 540 |
| Met Thr Val Val Ile Ala Gly Leu Ser Asn Val Tyr Thr His Tyr |     |     |
| 545   | 550 | 555 |
| Ile Thr Thr Tyr Glu Glu Tyr Gln Ala Gln Arg Tyr Glu Ala Ala |     |     |
| 560   | 565 | 570 |
| Ser Thr Ile Tyr Gly Pro His Thr Leu Ser Ala Tyr Ile Gln Leu |     |     |
| 575   | 580 | 585 |
| Phe Arg Asp Leu Ala Lys Ala Ile Ala Thr Asp Thr Val Ala Asn |     |     |
| 590   | 595 | 600 |
| Met Ser Ser Gly Pro Glu Pro Pro Phe Phe Lys Asn Leu Ile Ala |     |     |
| 605   | 610 | 615 |
| Ser Leu Ile Pro Asn Ile Ala Asp Arg Ala Pro Ile Gly Lys His |     |     |
| 620   | 625 | 630 |
| Phe Gly Asp Val Leu Gln Pro Ala Lys Pro Glu Tyr Arg Val Gly |     |     |
| 635   | 640 | 645 |
| Glu Val Val Glu Val Ile Phe Val Gly Ala Asn Pro Lys Asn Ser |     |     |
| 650   | 655 | 660 |
| Ala Glu Asn Gln Thr His Gln Thr Phe Leu Thr Val Glu Lys Tyr |     |     |
| 665   | 670 | 675 |
| Glu Asp Ser Val Ala Asp Trp Gln Ile Met Tyr Asn Asp Ala Ser |     |     |
| 680   | 685 | 690 |
| Trp Glu Thr Arg Phe Tyr Trp His Lys Gly Ile Leu Gly Leu Ser |     |     |
| 695   | 700 | 705 |
| Asn Ala Thr Ile Tyr Trp His Ile Pro Asp Thr Ala Tyr Pro Gly |     |     |
| 710   | 715 | 720 |
| Ile Tyr Arg Ile Arg Tyr Phe Gly His Asn Arg Lys Gln Glu Leu |     |     |
| 725   | 730 | 735 |

Leu Lys Pro Ala Val Ile Leu Ala Phe Glu Gly Ile Ser Ser Pro

740

745

750

Phe Glu Val Val Thr Thr

755

<210> 14

<211> 682

<212> PRT

<213> Mouse

<400> 14

Phe Ser Gly Tyr Tyr Ile Gly Val Gly Arg Ala Asp Cys Thr Gly

1

5

10

15

Gln Val Ser Asp Ile Asn Leu Met Gly Tyr Gly Lys Asn Gly Gln

20

25

30

Asn Ala Arg Gly Leu Leu Thr Arg Leu Phe Ser Arg Ala Phe Ile

35

40

45

Leu Ala Asp Pro Asp Gly Ser Asn Arg Met Ala Phe Val Ser Val

50

55

60

Glu Leu Cys Met Ile Ser Gln Arg Leu Arg Leu Glu Val Leu Lys

65

70

75

Arg Leu Glu Ser Lys Tyr Gly Ser Leu Tyr Arg Arg Asp Asn Val

80

85

90

Ile Leu Ser Ala Ile His Thr His Ser Gly Pro Ala Gly Phe Phe

95

100

105

Gln Tyr Thr Leu Tyr Ile Leu Ala Ser Glu Gly Phe Ser Asn Arg

110

115

120

Thr Phe Gln Tyr Ile Val Ser Gly Ile Met Lys Ser Ile Asp Ile

125

130

135

09/937521 1254660

|   |     |     |
|---|-----|-----|
| Ala His Thr Asn Leu Lys Pro Gly Lys Ile Phe Ile Asn Lys Gly |     |     |
| 140   | 145 | 150 |
| Asn Val Ala Asn Val Gln Ile Asn Arg Ser Pro Ser Ser Tyr Leu |     |     |
| 155   | 160 | 165 |
| Leu Asn Pro Gln Ser Glu Arg Ala Arg Tyr Ser Ser Asn Thr Asp |     |     |
| 170   | 175 | 180 |
| Lys Glu Met Leu Val Leu Lys Leu Val Asp Leu Asn Gly Glu Asp |     |     |
| 185   | 190 | 195 |
| Leu Gly Leu Ile Ser Trp Phe Ala Ile His Pro Val Ser Met Asn |     |     |
| 200   | 205 | 210 |
| Asn Ser Asn His Phe Val Asn Ser Asp Asn Met Gly Tyr Ala Ala |     |     |
| 215   | 220 | 225 |
| Tyr Leu Phe Glu Gln Glu Lys Asn Lys Gly Tyr Leu Pro Gly Gln |     |     |
| 230   | 235 | 240 |
| Gly Pro Phe Val Ala Gly Phe Ala Ser Ser Asn Leu Gly Asp Val |     |     |
| 245   | 250 | 255 |
| Ser Pro Asn Ile Leu Gly Pro His Cys Val Asn Thr Gly Glu Ser |     |     |
| 260   | 265 | 270 |
| Cys Asp Asn Asp Lys Ser Thr Cys Pro Asn Gly Gly Pro Ser Met |     |     |
| 275   | 280 | 285 |
| Cys Met Ala Ser Gly Pro Gly Gln Asp Met Phe Glu Ser Thr His |     |     |
| 290   | 295 | 300 |
| Ile Ile Gly Arg Ile Ile Tyr Gln Lys Ala Lys Glu Leu Tyr Ala |     |     |
| 305   | 310 | 315 |
| Ser Ala Ser Gln Glu Val Thr Gly Pro Val Leu Ala Ala His Gln |     |     |
| 320   | 325 | 330 |
| Trp Val Asn Met Thr Asp Val Ser Val Gln Leu Asn Ala Thr His |     |     |
| 335   | 340 | 345 |
| Thr Val Lys Thr Cys Lys Pro Ala Leu Gly Tyr Ser Phe Ala Ala |     |     |

0937521-09601

|   |     |     |
|---|-----|-----|
| 350   | 355 | 360 |
| Gly Thr Ile Asp Gly Val Ser Gly Leu Asn Ile Thr Gln Gly Thr |     |     |
| 365   | 370 | 375 |
| Thr Glu Gly Asp Pro Phe Trp Asp Thr Leu Arg Asp Gln Leu Leu |     |     |
| 380   | 385 | 390 |
| Gly Lys Pro Ser Glu Glu Ile Val Glu Cys Gln Lys Pro Lys Pro |     |     |
| 395   | 400 | 405 |
| Ile Leu Leu His Ser Gly Glu Leu Thr Ile Pro His Pro Trp Gln |     |     |
| 410   | 415 | 420 |
| Pro Asp Ile Val Asp Val Gln Ile Val Thr Val Gly Ser Leu Ala |     |     |
| 425   | 430 | 435 |
| Ile Ala Ala Ile Pro Gly Glu Leu Thr Thr Met Ser Gly Arg Arg |     |     |
| 440   | 445 | 450 |
| Phe Arg Glu Ala Ile Lys Lys Glu Phe Ala Leu Tyr Gly Met Lys |     |     |
| 455   | 460 | 465 |
| Asp Met Thr Val Val Ile Ala Gly Leu Ser Asn Val Tyr Thr His |     |     |
| 470   | 475 | 480 |
| Tyr Ile Thr Thr Tyr Glu Glu Tyr Gln Ala Gln Arg Tyr Glu Ala |     |     |
| 485   | 490 | 495 |
| Ala Ser Thr Ile Tyr Gly Pro His Thr Leu Ser Ala Tyr Ile Gln |     |     |
| 500   | 505 | 510 |
| Leu Phe Arg Asp Leu Ala Lys Ala Ile Ala Thr Asp Thr Val Ala |     |     |
| 515   | 520 | 525 |
| Asn Met Ser Ser Gly Pro Glu Pro Pro Phe Phe Lys Asn Leu Ile |     |     |
| 530   | 535 | 540 |
| Ala Ser Leu Ile Pro Asn Ile Ala Asp Arg Ala Pro Ile Gly Lys |     |     |
| 545   | 550 | 555 |
| His Phe Gly Asp Val Leu Gln Pro Ala Lys Pro Glu Tyr Arg Val |     |     |
| 560   | 565 | 570 |

09/937521

Gly Glu Val Val Glu Val Ile Phe Val Gly Ala Asn Pro Lys Asn  
 575 580 585  
 Ser Ala Glu Asn Gln Thr His Gln Thr Phe Leu Thr Val Glu Lys  
 590 595 600  
 Tyr Glu Asp Ser Val Ala Asp Trp Gln Ile Met Tyr Asn Asp Ala  
 605 610 615  
 Ser Trp Glu Thr Arg Phe Tyr Trp His Lys Gly Ile Leu Gly Leu  
 620 625 630  
 Ser Asn Ala Thr Ile Tyr Trp His Ile Pro Asp Thr Ala Tyr Pro  
 635 640 645  
 Gly Ile Tyr Arg Ile Arg Tyr Phe Gly His Asn Arg Lys Gln Glu  
 650 655 660  
 Leu Leu Lys Pro Ala Val Ile Leu Ala Phe Glu Gly Ile Ser Ser  
 665 670 675  
 Pro Phe Glu Val Val Thr Thr  
 680

&lt;210&gt; 15

&lt;211&gt; 2049

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 15

ttcagtggct actacattgg cgttgggaga gcggattgca caggacaagt gtcagatattc 60  
 aatttgatgg gctatggcaa aaatggccag aatgcacggg gtctcctcac caggctgttc 120  
 agccgtgctt ttatcttggc ggatccagat gggtc aaatc gaatggcatt tgtgagcgtg 180  
 gaactatgta tgatttccca acgactgagg ttggagggtcc tgaagagact agagagtaaa 240  
 tatggctctc tgtatgaag agacaatgtt atcctgagt ccattcacac acactctggc 300  
 ccagcagggt ttttccaata tacactctat atactcgcca gcgagggtt cagcaaccgg 360

acctttcagt acatagtctc tgggatcatg aagagcattg atatagctca cacaatctt 420  
 aaaccaggca aaatctttat caacaaagga aatgttgcta atgtgcagat caaccgaagc 480  
 ccctcctctt accttctgaa tccacagtca gagagagcaa ggtattcttc aaacacagac 540  
 aaggaaatgc tggctttgaa actggtggat ttgaatggag aagacttggg tcttatcagc 600  
 tggtttgcca tccaccccggt gagcatgaac aatagcaacc accttctgaa tagtgacaat 660  
 atgggctatg cggcttacct ttttgagcaa gaaaagaaca aaggctatct gcctggacag 720  
 ggaccgtttg tagcaggctt tgcttcatca aatctcggag acgtgtcacc caacattctt 780  
 ggcccgcatg gtgtcaacac aggggagtct tgtgacaacg acaagagcac ctgtccaac 840  
 ggtgggccta gcatgtgcat ggccagcgga cctggacaag acatgtttga gagcacacac 900  
 attataggac ggatcatcta tcagaaggcc aaggagctgt atgcctctgc ctcccaggag 960  
 gtgaccggcc cagtgttgc agctcaccag tgggtgaaca tgacagatgt gagcgtccag 1020  
 ctcaatgcca cacacacagt gaagacgtgt aaacctgccc tgggctacag ttttccgca 1080  
 ggcaaatg atggagtctt gggcctcaat attacacagg gaactacgga aggggatcca 1140  
 ttctgggaca ctcttcggga ccagctcttg ggaaaacat ctgaagagat tgtagagtgt 1200  
 cagaaaccca aaccaatcct gcttcacagt ggagagctga cgataccaca tccttggcaa 1260  
 ccagatattg ttgatgtca gattgttacc gttgggtcct tggccatagc tgctatccct 1320  
 ggggaattaa caaccatgtc gggacgaaga ttctgtgagg caattaaaaa agaatttgca 1380  
 ctttatggga tgaaggatat gaccgttgtt atcgcaggtc taagcaatgt ttataacat 1440  
 tacattacca catatgaaga ataccaggct cagcggtagc aggcagcatc tacaatctat 1500  
 ggaccacaca ccctgtctgc atacatcaa ctcttcagag accttgctaa ggcaattgct 1620  
 acggacacag tagccaacat gagcagtggc cccgagcctc cattcttcaa aaatctaata 1620  
 gcttcactta ttctaatat tgcggataga gcaccaattg gcaaacattt tggggatgtc 1680  
 ttgcagccag caaacctga atacagagtg ggagaagtgg ttgaagtat attttaggc 1740  
 gctaacccaa agaattcagc agagaaccag acccatcaaa ccttcctcac tgttgagaaa 1800  
 tacgaggact ctgtagctga ctggcagata atgtataacg atgcctcttg ggagacgagg 1860  
 ttttatiggc acaagggaat actgggtctg agcaatgcaa caatatactg gcatattcca 1920  
 gatactgcct accctggaat ctacagaata agatatttg gacacaatcg gaagcaggaa 1980  
 cttctgaaac ccgctgcat actagcattt gaaggaattt cttctccttt tgaagttgtc 2040  
 actacttag 2049

&lt;210&gt; 16

&lt;211&gt; 4835

&lt;212&gt; DNA

&lt;213&gt; Mouse

&lt;400&gt; 16

cctgcagcgg tgttctgaag agccgggcag aggatacaca agcatcccag caggcactct 60  
ggtttgcccg tgaacgatag atatgcgggg gtttgaatga gcagctgcag cagcggggtt 120  
gggtctgtac acatgagtgc tgggtgtgctt agaggcatcg ggtccctttc agctggagtt 180  
gcagtacttg tgagtgccat atttggaac acattgagaa atcagcctga gcaacctgca 240  
aggcacaagg cacaagattc tgcatggtaa tttgtctcc caggaggtga acatttggtt 300  
tgattaacag agtcagggtt gagatgccc gttgttcctc atcttggctc agaagaagca 360  
cctaggaata aaagctctaa gctggtatta agtagaatgg gcttaaagtc cactacagga 420  
aacaacagct agtgacagaa atggcaaagc gaaccttctc caccttggag gcatttcctca 480  
ttttccttct ggtaataatg acagtcatca cagtggccct tctcacctc ttgtttgta 540  
ccagtgggac cattgaaaac cacaagatt caggaaatca ctggttttca accactctgg 600  
gctccacgac aaccagccc cctccaatta cacagactcc aaacttcctt icatttcgga 660  
acttcagtgg ctactacatt ggcgttggga gacagattg cacaggacaa gtgtcagata 720  
tcaatttgat gggctatggc aaaaatggcc agaatgcacg ggtctcctc accaggctgt 780  
tcagccgtgc ttttatcttg gcggatccag atgggtcaaa tcgaatggca tttgtgagcg 840  
tggaactatg tatgatttcc caacgactga ggttggaggt cctgaagaga ctagagagta 900  
aatatggctc tctgtatcga agagacaatg ttatcctgag tgccattcac acacactctg 960  
gcccgagcagg gtttttccaa tatacactct atatactcg cagcgaggga ttcagcaacc 1020  
ggacctttca gtacatagtc tctgggatca tgaagagcat tgatatagct cacacaaatc 1080  
ttaaaccagg caaaatcttt atcaacaaag gaaatgttgc taatgtgcag atcaaccgaa 1140  
gcccctctc ttaacctctg aatccacagt cagagagagc aaggtattct tcaaacacag 1200  
acaaggaaat gctggtcttg aaactggtgg atttgaatgg agaagacttg ggtcttatca 1260  
gctggtttgc catccacccc gtgagcatga acaatagcaa ccactttgtg aatagtgaca 1320

atatgggcta tgcggcttac ctttttgagc aagaaaagaa caaaggctat ctgcctggac 1380  
 agggaccgtt tgtagcaggc ttgtctcat caaatctcgg agacgtgtca cccaacattc 1440  
 ttggcccgca ttgtgtcaac acaggggagt cttgtgacaa cgacaagagc acctgtccca 1500  
 acggtgggcc tagcatgtgc atggccagcg gacctggaca agacatgttt gagagcacac 1560  
 acattatagg acggatcatc tatcagaagg ccaaggagct gtatgcctct gcctcccagg 1620  
 aggtgaccgg ccagtgctt gcagctcacc agtgggtgaa catgacagat gtgagcgtcc 1680  
 agtcaatgc cacacacaca gtgaagacgt gtaaacctgc cctgggctac agttttgccg 1740  
 caggcacaat tgatggagtt tcgggcctca atattacaca gggaactacg gaaggggatc 1800  
 cattctggga cactcttcgg gaccagctct tgggaaaacc atctgaagag attgttagagt 1860  
 gtcagaaacc caaaccaatc ctgcttcaca gtggagagct gacgatacca catccttggc 1920  
 aaccagatat tgttgatgtt cagattgtta ccgttgggtc cttggccata gctgctatcc 1980  
 ctggggaatt aacaacatg tcgggacgaa gatttcgtga ggcaattaaa aaagaatttg 2040  
 cactttatgg gatgaaggat atgaccgttg ttatcgcagg tctaagcaat gttatacac 2100  
 attacattac cacatatgaa gaataccagg ctacagcgta cgaggcagca tctacaatct 2160  
 atggaccaca caccctgtct gcatacatcc aactcttcag agaccttgct aaggcaattg 2220  
 ctacggacac agtagccaac atgagcagtg gtcccagacc tccattcttc aaaaatctaa 2280  
 tagcttcact tattcctaatt attgcggata gaccaccaat tggcaaacat tttggggatg 2340  
 tcttgagcc agcaaacct gaatacagag tgggagaagt ggttgaagtt atattttag 2400  
 gcgctaacc aaagaattca gcagagaacc agacccatca aaccttcctc actgtggaga 2460  
 aatacgagga ctctgtagct gactggcaga taatgtataa cgatgcctcc tgggagacga 2520  
 ggttttatg gcacaaagga atactgggtc tgagcaatgc aacaatatac tggcatattc 2580  
 cagatactgc ctaccctgga atctacagaa taagatatit tggacacaat cggaagcagg 2640  
 aacttctgaa acccgctgtc atactagcat ttgaaggaat ttcttctcct tttgaagttg 2700  
 tcactactta gtgaaaagtt gacagatatt gaagaaaagc ttttctctgt gcacattata 2760  
 gagtgaattc acacaaatgt gaactgccag ttttaattct gtaattgtct ctgtttggg 2820  
 gacaggtcat ttattgctaa tgggacagag gtatgtgttt gtgtgtgtgt atgattatga 2880  
 gtatgcatgc taacaggaag agagagggag gagggagggg ggagggggag ggaggggaaga 2940  
 aaggagggag agagagagtg agagaatgag agagagagtg agagagaaag agttattagt 3000  
 gagcaagaga atatgagaga agggccactg acaaccaaatt acctgtgat ctttatccta 3060

0937521-254660

aagcatgatt ttccttgaag ctctgtgggt gttaagaga taattccctc taatatgaaa 3120  
tccctgaaat ataatgacag tatttgaaga tatgtgaata atgtttatcc tatttattta 3180  
tagacttact aaatgagaac actagagaac ttcttagaag tcctctagaa tgatacttga 3240  
ttttacagag aggaaaagga gctttgattc tctttagggt agaataaggt tagtatattt 3300  
ttccctagtc atatttacia aataccatgt aactttacta caaatatttg agcccagcta 3360  
aaatataccc agaaaattag cataccaggt ttgttttgggt ttattttggt ttgcatcca 3420  
aacaagcata gtccttctga taagtcactt tagaatggat ctgcctggct cagggttatt 3480  
gttcatgctc agatcatttc cgcaattacc tccagagtcc aactatgca atgtcacttg 3540  
cagtgccttg atttatgcct tgtattcctc aaagtgcct tatcctgcta agtcacacct 3600  
cttctccca gcatttactc taaatgattt ttaatgtttt cgccaatcaa atgtacctca 3660  
cattacaaag ctttgccttg aatgtagatt tttaaaacaa aagtgttaag gctggaaatg 3720  
tagttatcaa agaggaaggt ttaaattgat ctgttctttt atcagctact ccttccctca 3780  
tggtccctt gaatcactga atagttattt aaaccacat atccaatatg gtactcattc 3840  
ctgggtcttc acaattacag acatcatatc gaaatgattg ggctgacaat tcctttgaag 3900  
gacaaagtaa atatttaattg agaaatatag attctggaga ggcaattgaa aatcacaaat 3960  
gttacgcctc catttctgt ttccaggtt ggggtttctg atttggagg aaagcagccc 4020  
caaataattt taaatatga atctgaaaat aatgttttag aaattatgat ctgcacagtc 4080  
taattaatga gaattgtctg aaagtcctag ctgcatttaa aattatgtaa gttactaaa 4140  
gccaatttt gaacccagct cataatttg taggtaggta aaaagagcat tttaggagga 4200  
aaccgaactt catttcaaga ctgaatctgt tttaaaagaa caatagtgtt aaggtaaatc 4260  
ttcatttatt tccctatggt ttacctattt aaacatcgaa gattgaatca aaaggcacct 4320  
ggagcatatt ttgtaactc catttccac ttgtagttc tatggatgct aactgctgaa 4380  
gaataaactg atcgggattt tcaagggttg tgaacatgct tcctgatggg aataccgtat 4440  
taagtataaa ggttcaaaat agttgatctc aaaactatac acacacacac aatatatata 4500  
tatatacaca cacacacatg tacacacaca cacacatgca catacacatg gtattgttta 4560  
aaatttattt ctcatgactt agaacaatat aaggattata caaggattca ttcccacca 4620  
tcattctctc cagtgaagct ttctcaaag tctgagtagg agtttctctt ttctactgg 4680  
taactatccc acagtggcca ttacatcact agtaatcggt gtgcccagcc ctgcatggaa 4740  
ataaatcaca gaaacataat ttccagtag acttagtctc ttcaagcctg tgtgcttcta 4800

gtgtataaaa tctgtaaaaa aaaaaaaaaa aaaaa

4835

&lt;210&gt; 17

&lt;211&gt; 24

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Synthesized oligonucleotide for primer

&lt;400&gt; 17

gtttgagagc acacacatta tagg

24

&lt;210&gt; 18

&lt;211&gt; 24

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Synthesized oligonucleotide for primer

&lt;400&gt; 18

atattgaggc ccgaaactcc atca

24

T09260" T254660